

NONLERFNESS OF ARITHMETIC HYPERBOLIC MANIFOLD GROUPS

HONGBIN SUN

ABSTRACT. We will show that, for any noncompact arithmetic hyperbolic m -manifold with $m \geq 4$, and any compact arithmetic hyperbolic m -manifold with $m \geq 5$ which is not a 7-dimensional arithmetic hyperbolic manifold defined by octonions, its fundamental group is not LERF. The main ingredient in the proof is a study of certain graph of groups with hyperbolic 3-manifold groups being the vertex groups. We will also show that a compact irreducible 3-manifold with empty or tori boundary does not support a geometric structure if and only if its fundamental group is not LERF.

1. INTRODUCTION

For a group G and a subgroup $H < G$, we say that H is *separable* in G if for any $g \in G \setminus H$, there exists a finite index subgroup $G' < G$ such that $H < G'$ and $g \notin G'$. G is called *LERF* (locally extended residually finite) or *subgroup separable* if all finitely generated subgroups of G are separable.

The LERFness of a group is a property closely related with low dimensional topology, especially the virtual Haken conjecture (which is settled in [Ag2]). So in this paper, we are mostly interested in fundamental groups of some nice manifolds, and graph of groups constructed from these groups.

Among fundamental groups of low dimensional manifolds, it is known that the following groups are LERF: free groups ([Ha]), surface groups ([Sc]), Seifert manifold groups ([Sc]), hyperbolic 3-manifolds groups ([Ag2] and [Wi]); while the following groups are known to be nonLERF: the groups of nontrivial graph manifolds ([NW2]), the groups of fibered 3-manifolds whose monodromy is reducible and satisfies some further condition ([Li]).

In this paper, we give a few more examples of nonLERF groups arising from topology. These results imply that 3-manifolds with LERF fundamental groups support geometric structures, and hyperbolic manifolds with LERF fundamental groups seem to have dimension at most 3.

One main result of this paper is about high dimensional arithmetic hyperbolic manifolds (dimension ≥ 4). Comparing to the 3-dimensional case, there are much fewer examples of hyperbolic manifolds with dimension at least 4. Most examples of high dimensional hyperbolic manifolds are constructed by arithmetic methods, and some other examples are constructed by doing cut-and-paste surgery on these arithmetic examples. So the following results suggest that having nonLERF fundamental group is a general phenomenon in high dimensional hyperbolic world.

Date: August 18, 2016.

2010 Mathematics Subject Classification. 20E26, 57M05, 57M50, 22E40.

Key words and phrases. locally extended residually finite, graph of groups, hyperbolic 3-manifolds, arithmetic hyperbolic manifolds.

The author is partially supported by NSF Grant No. DMS-1510383.

Theorem 1.1. *Let M^m be an arithmetic hyperbolic manifold with $m \geq 5$ which is not a 7-dimensional arithmetic hyperbolic manifold defined by octonions, then its fundamental group is not LERF.*

Moreover, if M is closed, there exists a free product of closed surface groups and free groups as a nonseparable subgroup. If M is not closed, there exists either a free subgroup or a free product of closed surface groups and free groups as a nonseparable subgroup.

Comparing to Theorem 1.1, it is shown in [BHW] that all geometrically finite subgroups of standard arithmetic hyperbolic manifold groups are separable. It will be easy to see that the nonseparable subgroups constructed in the proof of Theorem 1.1 are not geometrically finite.

Theorem 1.1 does not cover the case of arithmetic hyperbolic 4-manifolds. By using a slightly different method, we can show that noncompact arithmetic hyperbolic manifolds with dimension at least 4 have nonLERF fundamental groups. Of course, the only case in Theorem 1.2 which is not covered by Theorem 1.1 is the 4-dimensional case.

Theorem 1.2. *Let M^m be a noncompact arithmetic hyperbolic m -manifold with $m \geq 4$, then $\pi_1(M)$ is not LERF.*

Moreover, there exist both a free subgroup and a closed surface subgroup as nonseparable subgroups.

Some examples of high dimensional nonarithmetic hyperbolic manifolds are constructed in [GPS], [Ag1] and [BT]. These examples are constructed by cutting arithmetic hyperbolic manifolds along codimension-1 totally geodesic submanifolds, then pasting along isometric boundary components. Since all these nonarithmetic hyperbolic manifolds contain codimension-1 arithmetic hyperbolic submanifolds, Theorem 1.1 implies Theorem 5.1, which claims that all the nonarithmetic examples in [GPS] and [BT] ([Ag1] only constructed 4-dimensional examples) with dimension ≥ 6 have nonLERF fundamental groups.

In Theorem 5.2, we also show that compact reflection hyperbolic manifolds with dimension ≥ 5 and noncompact reflection hyperbolic manifolds with dimension ≥ 4 have nonLERF fundamental groups.

Another main result in this paper concerns irreducible compact 3-manifolds with empty or tori boundary. Thurston's Geometrization Conjecture (confirmed by Perelman) implies that any irreducible compact 3-manifolds M with empty or tori boundary can be cut along a minimal set of incompressible tori, such that each component of the complement supports one of eight Thurston's geometries. If this set of incompressible tori is empty, we say that M is a geometric 3-manifold.

The following theorem implies that an irreducible compact 3-manifolds with empty or tori boundary is geometric if and only if its fundamental group is LERF. The author thinks that this result is very interesting, since it gives a surprising relation between geometric structures of 3-manifolds and the LERFness of 3-manifold groups. These two topics in 3-manifold topology are both very popular in the past twenty years. This result also confirms Conjecture 1.5 in [Li1].

Theorem 1.3. *For an irreducible compact orientable 3-manifold M with empty or tori boundary, M supports one of eight Thurston's geometries if and only if $\pi_1(M)$ is LERF.*

When $\pi_1(M)$ is not LERF, there exists a free subgroup as a nonseparable subgroup. If M is a closed mixed 3-manifold, there also exists a closed surface subgroup as a nonseparable subgroup.

To prove Theorem 1.3, we only need to prove non-geometric 3-manifolds have nonLERF fundamental groups. Our proof is similar to the construction in Section 8 of [Li1]. Actually, there was an error in [Li1], which affects some statements in [Li1], but does not affect the main theorem over there (see [Li2]). If there was not the error in [Li1], the author believes that Yi Liu would be able to prove Theorem 1.3, since all the tools were already available for him.

To prove Theorem 1.3, the main case we need to deal with is that M is a union of two geometric 3-manifolds along one torus, with one of them being hyperbolic. From the group theory point of view, this group is a $\mathbb{Z}^2$ -amalgamation of two LERF groups. So it is natural to consider a simpler case: a $\mathbb{Z}$ -amalgamation of two hyperbolic 3-manifold groups, i.e. the fundamental group of a union of two hyperbolic 3-manifolds along one essential circle.

There have been a lot of works on the LERFness of $\mathbb{Z}$ -amalgamated groups $A *_\mathbb{Z} B$, with both A and B being LERF. Here we list some of them. The first nonLERF example of $A *_\mathbb{Z} B$ was constructed in [Ri]. It has been shown that if both A and B are free groups ([BBS]), or if A is free, B is LERF and $\mathbb{Z} < A$ does not lie in a bigger cyclic subgroup ([Gi]), or if both A and B are surface groups ([Ni1]), then $A *_\mathbb{Z} B$ is LERF.

Here we give a family of nonLERF $\mathbb{Z}$ -amalgamations of 3-manifold groups.

Theorem 1.4. *Let M_1, M_2 be two finite volume hyperbolic 3-manifolds, and $i_k : S^1 \rightarrow M_k$, $k = 1, 2$ be two embedded essential circles, then the fundamental group of*

$$X = M_1 \cup_{S^1} M_2$$

is not LERF.

Moreover, if both M_1 and M_2 are not closed, there exists a free subgroup as a nonseparable subgroup; if at least one of M_k is closed, there exists a free product of closed surface groups and free groups as a nonseparable subgroup.

Theorem 1.4 is the main ingredient to prove Theorem 1.1. We will use the fact that arithmetic hyperbolic manifolds have a lot of totally geodesic submanifolds of any smaller dimension. If a arithmetic hyperbolic manifold has dimension at least 5, there are two totally geodesic 3-dimensional submanifolds intersecting along a closed geodesic, which gives a picture already addressed in Theorem 1.4.

In dimension 4, such a picture does not show up because of the dimension reason, so Theorem 1.4 does not help here. However, Theorem 1.3 implies that the double of any cusped hyperbolic 3-manifold has nonLERF fundamental group, and the groups of all noncompact arithmetic hyperbolic manifolds with dimension ≥ 4 contain such doubled manifold groups (by [LR]). So Theorem 1.2 is a consequence of Theorem 1.3.

The organization of this paper is as the following. In Section 2, we review some background on group theory, 3-manifold topology and arithmetic hyperbolic manifolds. In Section 3, we prove Theorem 1.3, which is enlightened by the construction in [Li1], with some modifications. In Section 4, we prove Theorem 1.4, whose proof is similar to the proof of Theorem 1.3, with some further modifications. In Section

5, we deduce Theorem 1.1 and Theorem 1.2 from Theorem 1.4 and Theorem 1.3 respectively. In Section 6, we ask some questions related to the results in this paper.

Acknowledgement. The author is grateful to Ian Agol for many valuable conversations, they are very helpful on various aspects of the development of this paper. The author thanks Yi Liu for communication and explanation on his results in [Li1], and thanks Alan Reid for his help on arithmetic hyperbolic manifolds. Part of the work in this paper is done during the author's visiting at the Institute for Advanced Study, so the author thanks for the hospitality of IAS.

2. PRELIMINARIES

In this section, we review some basic material on group theory, 3-manifold topology and arithmetic hyperbolic manifolds.

2.1. Locally extended residually finite. In this subsection, we review basic concept and properties on locally extended residually finite groups.

Definition 2.1. Let G be a group, and $H < G$ be a subgroup, we say that H is *separable* in G if for any $g \in G \setminus H$, there exists a finite index subgroup $G' < G$ such that $H < G'$ and $g \notin G'$.

An equivalent formulation is that H is separable in G if and only if H is a closed set under the profinite topology of G .

Definition 2.2. A group G is *LERF* (locally extended residually finite) or *subgroup separable* if all finitely generated subgroups of G are separable in G .

A basic property on LERFness of groups is that any subgroup of a LERF group is still LERF. This property is basic and well-known, and the proof is very simple. However, since we will use this property for many times in this paper, we give a proof here.

Lemma 2.3. *Let G be a group and $\Gamma < G$ be a subgroup. For a further subgroup $H < \Gamma$, if H is separable in G , then H is separable in Γ .*

In particular, if Γ is not LERF, then G is not LERF.

Proof. For any $\gamma \in \Gamma \setminus H \subset G \setminus H$. Since H is separable in G , there exists a finite index subgroup $G' < G$ such that $H < G'$ and $\gamma \notin G'$. Then $\Gamma' = G' \cap \Gamma$ is a finite index subgroup of Γ , with $H < G' \cap \Gamma = \Gamma'$ and $\gamma \notin G' \cap \Gamma = \Gamma'$. So H is also separable in Γ .

If Γ is not LERF, it contains a finitely generated subgroup H which is not separable in Γ . Then the previous paragraph implies that H is not separable in G . So G is not LERF. $\square$

In this paper, the main method to prove a group is not LERF is to find a descending tower of subgroups in it, until we get a subgroup which has a nice structure such that a geometric argument can be applied to prove its nonLERFness. Then we can apply Lemma 2.3.

2.2. Geometric decomposition of irreducible 3-manifolds. In this paper, we assume all manifolds are connected and oriented, all 3-manifolds are compact and have empty or tori boundary. For any noncompact finite volume hyperbolic manifold M , we can add the ideal boundary to the cusps of M , which has a Euclidean structure.

Let M be an irreducible 3-manifold with empty or tori boundary. By the geometrization of 3-manifolds, which is achieved by Perelman and Thurston, exactly one of the following hold.

- M is geometric, i.e. M supports one of the following eight geometries: $\mathbb{E}^3$, $\mathbb{S}^3$, $\mathbb{S}^2 \times \mathbb{E}^1$, $\mathbb{H}^2 \times \mathbb{E}^1$, Nil, Sol, $\widetilde{\text{PSL}_2(\mathbb{R})}$, $\mathbb{H}^3$.
- There is a nonempty minimal union $\mathcal{T}_M \subset M$ of disjoint essential tori and Klein bottles, unique up to isotopy, such that each component of $M \setminus \mathcal{T}_M$ is either Seifert fibered or atoroidal. In the Seifert fibered case, this piece supports the $\mathbb{H}^2 \times \mathbb{E}^1$ -geometry and the $\widetilde{\text{PSL}_2(\mathbb{R})}$ -geometry; in the atoroidal case, this piece supports the $\mathbb{H}^3$ -geometry.

If M has nontrivial geometric decomposition as in the second case, we say that M is a *non-geometric 3-manifold*, and call components of $M \setminus \mathcal{T}_M$ *Seifert pieces* or *hyperbolic pieces*, according to their geometry. If the components of $M \setminus \mathcal{T}_M$ are all Seifert pieces, M is called a *graph manifold*. Otherwise, M contains a hyperbolic piece, and it is called a *mixed manifold*. Since we only care about virtual properties of 3-manifolds, we can pass to a double cover and assume all components of $\mathcal{T}_M$ are tori.

The geometric decomposition is very closely related to, but slightly different from the more traditional JSJ decomposition. Since these two decompositions agree with each other on some finite cover of M , and we are studying virtual properties, we will not make much difference between them.

2.3. Fibered structures of 3-manifolds. In the construction of nonseparable subgroups in Theorem 1.3 and Theorem 1.4, all these subgroups have graph of group structures, and the vertex groups are groups of fibered surfaces in 3-manifolds. So we briefly review the theory of Thurston norm and its relation with fibered structures of 3-manifolds.

If a 3-manifold M has a surface bundle over circle structure, and $b_1(M) > 1$, then M has infinitely many different surface bundle structures. (Actually this works for all dimensions.) These fibered structures of the 3-manifold M are organized by the Thurston norm on $H_2(M, \partial M; \mathbb{R}) (\cong H^1(M; \mathbb{R})$ by duality) defined in [Th].

For any $\alpha \in H_2(M, \partial M; \mathbb{Z}) (\cong H^1(M; \mathbb{Z}))$, its Thurston norm is defined by:

$$\|\alpha\| = \inf \{ |\chi(T_0)| \mid (T, \partial T) \subset (M, \partial M) \text{ represents } \alpha \},$$

where $T_0 \subset T$ excludes S^2 and D^2 components of T . In [Th], it is shown that the norm can be extended to $H_2(M, \partial M; \mathbb{R})$ homogeneously and continuously, and the Thurston norm unit ball is a polyhedron with faces dual with elements in $H_1(M; \mathbb{Z})/\text{Tor}$. For a general 3-manifold, the Thurston norm is only a semi-norm, and it is a genuine norm for finite volume hyperbolic 3-manifolds.

For a top dimensional open face F of the Thurston norm unit ball, let C be the open cone over F . In [Th], Thurston showed that an integer point $\alpha \in H_2(M, \partial M; \mathbb{R})$ corresponds to a surface bundle structure of M if and only if α is contained in an open cone C as above and all integer points in C correspond

to surface bundle structures of M . In this case, the corresponding face F is called a *fibred face*, and C is called a *fibred cone*. For any point (possibly not an integer point) in a fibred cone, we call it a *fibred class*.

Thurston's theorem implies that the set of fibred classes of M is an open subset of $H_2(M, \partial M; \mathbb{R})$. In particular, if we start with a fibred class $\alpha \in H_2(M, \partial M; \mathbb{R})$, then for any $\beta \in H_2(M, \partial M; \mathbb{R})$, there exists $\epsilon > 0$, such that $\alpha + c\beta \in H_2(M, \partial M; \mathbb{R})$ is a fibred class for any $c \in (-\epsilon, \epsilon)$.

2.4. Virtual retractions of hyperbolic 3-manifold groups. Suppose we are given a fibred class $\alpha \in H_2(M, \partial M; \mathbb{R})$, then we may want to perturb this fibred class such that the perturbation has some desired property. To make sure the desired perturbation exists, we need some group theoretical property of hyperbolic 3-manifold groups.

The property we need is called virtual retract property, which is defined as the following.

Definition 2.4. For a group G and a (infinite index) subgroup $H < G$, we say that H is a *virtual retraction* of G if there exists a finite index subgroup $G' < G$ and a homomorphism $\phi : G' \rightarrow H$, such that $H < G'$ and $\phi|_H = id_H$.

For a finite volume hyperbolic 3-manifold M , the following dichotomy for finitely generated subgroups $H < \pi_1(M)$ holds.

- (1) H is a geometrically finite subgroup of $\pi_1(M)$, from the Kleinian group point of view. Equivalently, H is (relatively) quasiconvex in the (relative) hyperbolic group $\pi_1(M)$, from the geometric group theory point of view.
- (2) H is a geometrically infinite subgroup of $\pi_1(M)$. In this case, H is a virtual fibred surface subgroup of M .

Here we do not give the definition of geometrically finite and geometrically infinite subgroups, the readers only need to know that if H is not a virtual fibred surface subgroup, then it is a geometrically finite subgroup.

In [CDW], it is shown that (relatively) quasiconvex subgroups of virtually compact special (relative) hyperbolic groups are virtual retractions. The celebrated virtual compact special theorem of Wise ([Wi] for cusped case) and Agol ([Ag2] for closed case) implies that the groups of finite volume hyperbolic 3-manifolds are virtually compact special. These two results together give us the following theorem.

Theorem 2.5. *Let M be a finite volume hyperbolic 3-manifold, $H < \pi_1(M)$ be a geometrically finite subgroup (i.e. H is not a virtual fibred surface subgroup), then H is a virtual retraction of $\pi_1(M)$.*

2.5. Arithmetic hyperbolic manifolds. In this subsection, we briefly review the definition of (standard) arithmetic hyperbolic manifolds. Most of the material can be found in Chapter 6 of [VS].

Recall that the hyperboloid model of $\mathbb{H}^n$ is given as the following. Equip $\mathbb{R}^{n+1}$ with a bilinear form $B : \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ defined by

$$B((x_1, \dots, x_n, x_{n+1}), (y_1, \dots, y_n, y_{n+1})) = x_1 y_1 + \dots + x_n y_n - x_{n+1} y_{n+1}.$$

Then the hyperbolic space $\mathbb{H}^n$ is identified with

$$I^n = \{\vec{x} = (x_1, \dots, x_n, x_{n+1}) \mid B(\vec{x}, \vec{x}) = -1, x_{n+1} > 0\}.$$

The hyperbolic metric is given by the restriction of $B(\cdot, \cdot)$ on the tangent space of I^n .

The isometry group of $\mathbb{H}^n$ consists of all linear transformations of $\mathbb{R}^{n+1}$ which preserve $B(\cdot, \cdot)$ and fixe I^n . Let $J = \text{diag}(1, \dots, 1, -1)$ be the $(n+1) \times (n+1)$ matrix defining the bilinear form $B(\cdot, \cdot)$, then the isometry group of $\mathbb{H}^n$ is given by

$$\text{Isom}(\mathbb{H}) \cong PO(n, 1; \mathbb{R}) = \{X \in GL(n+1, \mathbb{R}) \mid X^t J X = J\} / (X \sim -X).$$

The orientation preserving isometry group of $\mathbb{H}^n$ is given by

$$\text{Isom}_+(\mathbb{H}^n) \cong SO_0(n, 1; \mathbb{R}),$$

which is the component of

$$SO(n, 1; \mathbb{R}) = \{X \in SL(n+1, \mathbb{R}) \mid X^t J X = J\}$$

containing the identity matrix.

Now we give the definition of *standard arithmetic hyperbolic manifolds*, which are also called *arithmetic hyperbolic manifolds of simplest type*.

Let $K \subset \mathbb{R}$ be a totally real number field, and $\sigma_1 = \text{id}, \sigma_2, \dots, \sigma_k$ be all the embeddings of K into $\mathbb{R}$. Let

$$f(x) = \sum_{i,j=1}^{n+1} a_{ij} x_i x_j, \quad a_{ij} = a_{ji} \in K$$

be a nondegenerate quadratic form defined over K with negative inertia index 1 (as a quadratic form over $\mathbb{R}$). Suppose that for any $l > 1$, the quadratic form

$$f^{\sigma_l}(x) = \sum_{i,j=1}^{n+1} \sigma_l(a_{ij}) x_i x_j$$

is positive definite, then the information of K and f can be used to define an arithmetic hyperbolic group.

Let $\mathcal{O}_K$ be the ring of algebraic integers in K , and A be the $(n+1) \times (n+1)$ matrix defining f . Since the negative inertia index of A is 1, the *special orthogonal group of f* :

$$SO(f; \mathbb{R}) = \{X \in SL(n+1, \mathbb{R}) \mid X^t A X = A\}$$

is conjugate to $SO(n, 1; \mathbb{R})$ by a matrix P (satisfying $P^t A P = J$). $SO(f; \mathbb{R})$ has two components, and let $SO_0(f; \mathbb{R})$ be the component containing the identity matrix.

Then we can form the set of algebraic integer points

$$SO(f; \mathcal{O}_K) = \{X \in SL(n+1, \mathcal{O}_K) \mid X^t A X = A\} \subset SO(f; \mathbb{R})$$

in $SO(f; \mathbb{R})$. The theory of arithmetic groups shows that

$$SO_0(f; \mathcal{O}_K) = SO(f; \mathcal{O}_K) \cap SO_0(f; \mathbb{R})$$

is conjugate to a lattice of $\text{Isom}_+(\mathbb{H}^n)$ (by the matrix P), i.e. it has finite volume quotient. For simplicity, we abuse notation and still use $SO_0(f; \mathcal{O}_K)$ to denote its P -conjugation in $SO_0(n, 1; \mathbb{R}) \cong \text{Isom}_+(\mathbb{H}^n)$.

Here $SO_0(f; \mathcal{O}_K) \subset \text{Isom}_+(\mathbb{H}^n)$ is called the *arithmetic group* defined by number field K and quadratic form f , and $\mathbb{H}^n / SO_0(f; \mathcal{O}_K)$ is a finite volume hyperbolic arithmetic orbifold. A hyperbolic n -manifold (orbifold) M is called a *standard arithmetic hyperbolic manifold (orbifold)* if M is commensurable with $\mathbb{H}^n / SO_0(f; \mathcal{O}_K)$ for some K and f .

The arithmetic orbifold $\mathbb{H}^n/SO_0(f; \mathcal{O}_K)$ is noncompact if and only if $f(\vec{x}) = 0$ has a nontrivial solution $\vec{x} \in K^{n+1}$. This happens only if $K = \mathbb{Q}$ (thus $\mathcal{O}_K = \mathbb{Z}$). When $n \geq 4$, $\mathbb{H}^n/SO_0(f; \mathcal{O}_K)$ is noncompact if and only if $K = \mathbb{Q}$.

For this paper, the most important property of standard arithmetic hyperbolic manifolds is that they contain a lot of finite volume hyperbolic 3-manifolds as totally geodesic submanifolds. This can be done by diagonalizing the matrix A and taking an indefinite 4×4 submatrix.

The above method of quadratic forms over number fields gives all even dimensional arithmetic hyperbolic manifolds (orbifolds). In odd dimension, there is another family of arithmetic hyperbolic manifolds (orbifolds), which are defined by (skew-Hermitian) quadratic forms over quaternion algebras. We do not give the definition of this family here. The readers can find a detailed definition in [LM].

This family of arithmetic hyperbolic manifolds defined by quaternions also have a lot of finite volume hyperbolic 3-manifolds as totally geodesic submanifolds. This can be done by diagonalizing the quadratic form over quaternions and taking a 2×2 submatrix. This fact is also used in [Ka].

In dimension 7, there is another way to construct arithmetic hyperbolic manifolds by using octonions. They are sporadic examples, and the author does not know whether these manifolds have totally geodesic (or π_1 -injective) 3-dimensional submanifolds. All the examples in this family are compact manifolds.

3. NONLERFNESS OF NON-GEOMETRIC 3-MANIFOLD GROUPS

In this section, we prove that the groups of non-geometric 3-manifolds are not LERF. The construction of nonseparable (surface) subgroups is enlightened by the construction in [Li1] (and also [RW]). The proof of nonseparability is essentially a computation of the spirality character defined in [Li1], then apply Theorem 1.1 of [Li1]. The difference is that we do some proper modifications of the construction in [Li1], and give an elementary proof of the nonseparability without using the spirality character explicitly. This proof also works for 3-manifolds with nonempty tori boundary.

3.1. Finite semicovers of non-geometric 3-manifolds. We first review the notion of finite semicovers of nongeometric 3-manifolds, which was introduced in [PW2].

Definition 3.1. Let M be a nongeometric 3-manifold with tori or empty boundary. A *finite semicover* of M is a compact 3-manifold N and a local embedding $f : N \rightarrow M$, such that its restriction on each boundary component of N is a finite cover to a decomposition torus or a boundary component of M .

For a finite semicover $f : N \rightarrow M$, the decomposition tori of N is the preimage of $\mathcal{T}_M$, and the restriction of f on each geometric piece of N is a finite cover to the corresponding geometric piece of M .

One important property of finite semicovers is given by the following lemma in [Li1].

Lemma 3.2. *If N is a connected finite semicover of a nongeometric 3-manifold M with empty or tori boundary, then N has an embedded lifting in a finite cover*

of M . In fact, the semi-covering $N \rightarrow M$ is π_1 -injective and $\pi_1(N)$ is separable in $\pi_1(M)$.

Remark 3.3. In [Li1], this lemma is stated only for closed irreducible nongeometric 3-manifolds, but it clearly also holds for irreducible nongeometric 3-manifolds with nonempty boundary. This is because that we can get a closed manifold by pasting cusped hyperbolic 3-manifolds, e.g. figure eight knot complements, to M along the boundary, then apply the closed manifold version of Lemma 3.2 and Lemma 2.3.

3.2. Reduction to non-geometric 3-manifolds with very simple dual graph. To prove Theorem 1.3, we reduce to the case that the dual graph of M has two vertices, two edges, and M has at least one hyperbolic piece.

Let M be an irreducible nongeometric 3-manifold with tori or empty boundary. It is known that all graph manifolds have nonLERF fundamental groups ([NW2]), so we can assume that M has at least one hyperbolic piece, i.e. M is a mixed 3-manifold.

The dual graph of M is a graph with vertices corresponding to geometric pieces of M and edges corresponding to decomposition tori. The following lemma is the first step of our reduction of 3-manifolds, which reduces the nonLERFness proof to a very simple case: the dual graph of M has only two vertices and one edge.

Lemma 3.4. *Let M be a mixed 3-manifold, then there exists a 3-manifold $N = N_1 \cup_T N_2$ such that the following hold.*

- (1) N_1 is a cusped hyperbolic 3-manifold, and N_2 is a geometric 3-manifold.
- (2) $N_1 \cap N_2 = T$ is a single torus, and $N = N_1 \cup_T N_2$ is a fibered 3-manifold.
- (3) N is a finite semicover of M , so $\pi_1(N)$ is a subgroup of $\pi_1(M)$.

Proof. By [PW1], we can find a finite cover of M such that it is a fibered 3-manifold, and still denote it by M .

We first suppose that M has at least two geometric pieces. Take any hyperbolic piece N_1 , and take another (distinct) geometric piece N_2 adjacent to N_1 . It is possible that $N_1 \cap N_2$ consists of more than one tori, and let T be one of them. Cut M along all the decomposition tori in $\mathcal{T}_M$ except T , then the component containing N_1 and N_2 is the desired N , which is clearly a finite semicover of M .

The fibered structure on M induces a fibered structure on N , since fibered structures of 3-manifolds are compatible with the geometric decomposition. It is easy to see all the other desired conditions hold for N .

If M has only one geometric piece N_1 , since the geometric decomposition is nontrivial, there is a decomposition torus T which is adjacent to N_1 on both sides. Then we take a double cover of M along T , and reduce it to the previous case. $\square$

By Lemma 2.3, to prove the nonLERFness of mixed 3-manifold groups, we only need to consider the case $M = M_1 \cup_T M_2$ as in Lemma 3.4 (we use M and M_i instead of N and N_i since we will do some further constructions). The dual graph of M has two vertices and one edge, which is not our desired model for constructing nonseparable subgroups. Actually, we need a cycle in the dual graph of the 3-manifold. So we use the following lemma to pass to a further finite semicover, such that its dual graph consists of two vertices and two edges connecting these two vertices.

Lemma 3.5. *Let $M = M_1 \cup_T M_2$ be a 3-manifold satisfying the conclusion of Lemma 3.4, then there exists a 3-manifold $N = N_1 \cup_{T \cup T'} N_2$ with nonempty boundary such that the following hold.*

- (1) N_1 is a cusped hyperbolic 3-manifold, and N_2 is a geometric 3-manifold.
- (2) $N_1 \cap N_2 = T \cup T'$ is a union of two tori, and $N = N_1 \cup_{T \cup T'} N_2$ is a fibered 3-manifold.
- (3) The homomorphism $H_1(T \cup T'; \mathbb{Z}) \rightarrow H_1(N_1; \mathbb{Z})$ induced by inclusion is injective.
- (4) N is a finite semicover of M , so $\pi_1(N)$ is a subgroup of $\pi_1(M)$.
- (5) There exists a fibered surface S in N , which is a union of two subsurfaces $S = S_1 \cup_{c \cup c'} S_2$, such that $S_i = S \cap N_i$ is connected, and each of $c = S \cap T$ and $c' = S \cap T'$ is a circle.

Proof. Claim. There exists a 3-manifold $N = N_1 \cup_{T \cup T'} N_2$ satisfying conditions (1)-(4).

We first give the proof of this claim.

Fix a base point of M_1 on T . For $\mathbb{Z}^2 \cong \pi_1(T) < \pi_1(M_1) < \text{Isom}_+(\mathbb{H}^3)$, take any $g \in \pi_1(M_1)$ which moves the fixed point of $\pi_1(T)$ on S_∞^2 to a different point. By the combination theorem of Kleinian groups, for large enough integer k , the subgroup of $\pi_1(M_1)$ generated by $\pi_1(T)$ and $g^k \pi_1(T) g^{-k}$ is the free product of these two subgroups, i.e. isomorphic to $\mathbb{Z}^2 * \mathbb{Z}^2$, and we denote it by H .

Since $H < \pi_1(M_1)$ is not a surface subgroup, it is a geometrically finite subgroup. By Theorem 2.5, we can find a finite cover N_1 of M_1 , such that $H < \pi_1(N_1)$ and there exists a retraction homomorphism $\pi_1(N_1) \rightarrow H$. Since hyperbolic 3-manifolds have LERF fundamental groups, by passing to a further finite cover (still denoted by N_1), we can assume that $g^k \notin \pi_1(N_1)$, and N_1 has at least three boundary components.

Since $g^k \notin \pi_1(N_1)$, any embedded arc γ in N_1 (starting from the lifted base point) corresponding to g^k connects two different boundary components of N_1 , and we denote them by T_1 and T'_1 . Note that the covering map $N_1 \rightarrow M_1$ maps both T_1 and T'_1 to T by homeomorphisms. Then $H < \pi_1(N_1)$ corresponds to the fundamental group of the union of T_1 , T'_1 and γ . Since $H = \pi_1(T_1 \cup T'_1 \cup \gamma)$ is a retraction of $\pi_1(N_1)$, $H_1(T_1 \cup T'_1 \cup \gamma; \mathbb{Z}) = H_1(T_1 \cup T'_1; \mathbb{Z})$ is a retraction of $H_1(N_1; \mathbb{Z})$. So condition (3) holds for N_1 .

If M_2 is a cusped hyperbolic 3-manifold, by doing the same construction for M_2 , we get a finite cover $N_2 \rightarrow M_2$ such that two boundary components T_2 and T'_2 of N_2 are mapped to T by homeomorphisms. By identifying T_1 with T_2 and T'_1 with T'_2 , we get a semifinite cover $N = N_1 \cup_{T \cup T'} N_2$ of M satisfying conditions (1)-(4). Here we use T to denote the image of T_1 and T_2 , and use T' to denote the image of T'_1 and T'_2 .

If M_2 is a Seifert fibered space, before doing the above construction for M_1 , we first do the following preparation. Since M is a fibered 3-manifold, we have $M = S \times I / \phi$, where $\phi : S \rightarrow S$ is a reducible homeomorphism on a surface S . By taking some finite cyclic cover M' of M along S , we can assume that M' has two adjacent geometric pieces, such that one of them is a cusped hyperbolic 3-manifold, and another one is $\Sigma \times S^1$ with $\chi(\Sigma) < 0$.

We take the union of these two adjacent pieces along a torus and get our new $M = M_1 \cup_T M_2$ with $M_2 = \Sigma \times S^1$. Then we do the same construction for M_1

as above to get a finite cover N_1 . For M_2 , let c be the boundary component of Σ corresponding to the boundary component T of M_2 . Since $\chi(\Sigma) < 0$, there exists a finite cover $\Sigma' \rightarrow \Sigma$ such that Σ' has two boundary components c_2 and c'_2 which are mapped to c by homeomorphisms.

Then $N_2 = \Sigma' \times S^1$ is a finite cover of M_2 . Let T_2 and T'_2 be the boundary components of N_2 corresponding to $c_2 \times S^1$ and $c'_2 \times S^1$ respectively, then they are both mapped to T by homeomorphisms. We can paste N_1 and N_2 together to get the desired finite semicover $N = N_1 \cup_{T \cup T'} N_2$.

This finishes the proof of the claim.

Now $N = N_1 \cup_{T \cup T'} N_2$ satisfies conditions (1)-(4), so we need to work on condition (5).

First take an arbitrary fibered class $\alpha \in H^1(N; \mathbb{Z})$. The corresponding fibered surface S might be more complicated than what we want in condition (5), since $S \cap N_i$, $S \cap T$ and $S \cap T'$ may not be connected.

Write N as $N = S \times I / \phi$, where $\phi : S \rightarrow S$ is a reducible self-homeomorphism of S . Let $\mathcal{C}$ be the set of reduced circles such that $\phi| : S \setminus \mathcal{C} \rightarrow S \setminus \mathcal{C}$ is either pseudo-Anosov or periodic on each ϕ -component.

We first suppose that there are two components S_1 and S_2 of $S \setminus \mathcal{C}$ such that $S_i \subset N_i$, and $S_1 \cap S_2$ contains two circles c and c' with $c \subset T$ and $c' \subset T'$. Take a positive integer k , such that ϕ^k preserves each component of $S \setminus \mathcal{C}$ and each component of $\mathcal{C}$. In this case $N' = ((S_1 \cup_{c \cup c'} S_2) \times I) / \phi^k$ is a finite semi-cover of N . Let $\bar{N}_i = S_i \times I / \phi^k$, and $\bar{T}$ and $\bar{T}'$ be the components of $\partial \bar{N}_1$ (also $\partial \bar{N}_2$) containing c and c' respectively. Then it is easy to check that $\bar{N} = \bar{N}_1 \cup_{\bar{T} \cup \bar{T}'} \bar{N}_2$ satisfies all the desired conditions.

If there are not such two components of $S \setminus \mathcal{C}$, we need to modify the fibered surface S . Take a tubular neighborhood $N(T_1)$ of T_1 in N_1 , and give it a coordinate by $N(T_1) = T_1 \times I = (S^1 \times I) \times S^1$ such that

$$S \cap N(T_1) = (\{a_1, a_2, \dots, a_k\} \times I) \times S^1,$$

with $a_1, \dots, a_k$ following a cyclic order on S^1 .

The fibered structure on $N(T_1)$ is given by a fibered structure of $S^1 \times I$, and then cross with S^1 . For any integer j , we change the fibered structure on $N(T_1)$ by taking a new fibered structure on $S^1 \times I$ as the following, then cross with S^1 . The new fibered structure on $S^1 \times I$ is given by a union of disjoint embedded arcs $I_i \subset S^1 \times I$, such that I_i connects $(a_i, 0)$ to $(a_{i+j}, 1)$ (modulo k), here $i = 1, 2, \dots, k$. This fibered structure on $N(T_1)$ can be pasted with the original fibered structure of $N \setminus N(T_1)$ to get a new fibered structure of N .

If we start from one component $S_1 \subset S \cap N_1$, take any component $S_2 \subset S \cap N_2$ such that $S_1 \cap S_2 \cap T' \neq \emptyset$. Then $S_1 \cap T_1$ and $S_2 \cap T_2$ are two families of parallel circles on T , but they may not have circles being identified with each other. Then we can use the above modification of the fibered structure and choose a proper j , such that the new fibered surface satisfies the assumption of the previous case. $\square$

Actually, condition (5) is not really necessary in the proof of Theorem 1.3, but it will make the immersed π_1 -injective surface constructed in Proposition 3.6 in a simple shape.

3.3. Construction of nonseparable surface subgroups. In this subsection, we will construct a π_1 -injective properly immersed surface in the 3-manifold $N =$

$N_1 \cup_{T \cup T'} N_2$ constructed in Lemma 3.5, then prove that this surface subgroup is not separable in $\pi_1(N)$.

The following proposition constructs a π_1 -injective properly immersed surface in N , which is our candidate of a nonseparable surface subgroup. Readers can compare this construction with the construction in Section 8 of [Li1].

Proposition 3.6. *For the 3-manifold $N = N_1 \cup_{T \cup T'} N_2$ and fibered surface $S = S_1 \cup_{c \cup c'} S_2$ constructed in Lemma 3.5, there exists a π_1 -injective properly immersed surface $i : \Sigma \looparrowright N$ such that the following hold.*

- (1) Σ is a union of connected surfaces as $\Sigma = (\Sigma_{1,1} \cup \Sigma_{1,2}) \cup (\cup_{k=1}^{2n} \Sigma_{2,k})$, with $i(\Sigma_{1,j}) \subset N_1$ and $i(\Sigma_{2,k}) \subset N_2$.
- (2) The restriction of i on $\Sigma_{1,j}$ and $\Sigma_{2,k}$ are all embeddings, and the images are fibered surfaces of N_1 and N_2 respectively.
- (3) Each $\Sigma_{2,k}$ is a copy of S_2 in N_2 . So $\Sigma_{2,k}$ intersects with both T and T' along exactly one circle.
- (4) $\Sigma_{1,1} \cap \Sigma_{2,1}$ consists of two circles s and s' , with $i(s) \subset T$ and $i(s') \subset T'$.
- (5) $\Sigma_{1,1} \cap T$ consists of A parallel copies of c , and $\Sigma_{1,1} \cap T'$ consists of B parallel copies of c' , with $A \neq B$.

Proof. When we cut N along $T \cup T'$ and cut S along $c \cup c'$, we will use T_i and T'_i to denote the copies of T and T' in N_i respectively, and use c_i and c'_i to denote the copies of c and c' in S_i respectively.

Let $\alpha \in H^1(N; \mathbb{Z})$ be the fibered class dual to S , and let $\alpha_1 = \alpha|_{N_1}$. Then $\alpha_1|_{T_1}$ is dual to $c_1 \subset T_1$, and $\alpha_1|_{T'_1}$ is dual to $c'_1 \subset T'_1$.

Since $H_1(T_1 \cup T'_1; \mathbb{Z}) \rightarrow H_1(N_1; \mathbb{Z})$ is injective, there exists a direct summand $A < H_1(N_1; \mathbb{Z})$ such that $A \cong \mathbb{Z}^4$ and $H_1(T_1 \cup T'_1; \mathbb{Z}) < A$. Since $\mathbb{Z}^4 \cong H_1(T_1 \cup T'_1; \mathbb{Z}) < A \cong \mathbb{Z}^4$ is a finite index subgroup, there exists a homomorphism $\tau : A \rightarrow \mathbb{Z}$ such that $\tau|_{H_1(T_1; \mathbb{Z})}$ is dual to a positive multiple of c_1 , and $\tau|_{H_1(T'_1; \mathbb{Z})} = 0$.

Let $\phi : H_1(N_1; \mathbb{Z}) \rightarrow A$ be a retraction given by the direct summand, then we get a cohomology class $\beta \in H^1(N_1; \mathbb{Z})$ defined by $\tau \circ \phi : H_1(N_1; \mathbb{Z}) \rightarrow \mathbb{Z}$. By the construction of τ , $\beta|_{T_1} = l\alpha_1|_{T_1}$ for some $l \in \mathbb{Z}_+$ and $\beta|_{T'_1} = 0$.

By the theory of Thurston norm, for large enough $n \in \mathbb{Z}_+$, $\alpha_{1,1} = n\alpha_1 + \beta$ and $\alpha_{1,2} = n\alpha_1 - \beta$ are both fibered classes in $H^1(N_1; \mathbb{Z})$. Here we can also assume that $n > l$ and $\gcd(n, l) = 1$.

Since $\alpha_{1,1}|_{T_1}$ is dual to $n + l$ copies of c_1 , $\alpha_{1,1}|_{T'_1}$ is dual to n copies of c'_1 , and $\gcd(n, l) = 1$, $\alpha_{1,1} \in H^1(N_1; \mathbb{Z})$ is a primitive class. Similarly, $\alpha_{1,2} \in H^1(N_1; \mathbb{Z})$ is also primitive.

Let $\Sigma_{1,1} \subset N_1$ be the connected fibered surface dual to $\alpha_{1,1} \in H^1(N_1; \mathbb{Z})$ and $\Sigma_{1,2} \subset N_1$ be the connected fibered surface dual to $\alpha_{1,2}$. Then $\Sigma_{1,1} \cap T_1$ consists of $A = n + l$ copies of c_1 (as oriented curves), $\Sigma_{1,1} \cap T'_1$ consists of $B = n$ copies of c'_1 ; $\Sigma_{1,2} \cap T_1$ consists of $n - l$ copies of c_1 , $\Sigma_{1,2} \cap T'_1$ consists of n copies of c'_1 . So $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap T_1$ and $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap T'_1$ consist of $2n$ (oriented) copies of c_1 and c'_1 respectively.

Note that both $S_2 \cap T_2$ and $S_2 \cap T'_2$ are exactly one (oriented) copy of c_2 and c'_2 respectively. If we take $2n$ copies of S_2 in N_2 , and denote them by $\Sigma_{2,k}$, with $k = 1, 2, \dots, 2n$, then we can identify parallel circles in $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap T_1$ with $(\cup_{k=1}^{2n} \Sigma_{2,k}) \cap T_2$ on T , and identify parallel circles in $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap T'_1$ with $(\cup_{k=1}^{2n} \Sigma_{2,k}) \cap T'_2$ on T' to get an immersed surface Σ . When we do the identification, we first identify one circle in $\Sigma_{1,1} \cap T_1$ with the circle in $\Sigma_{2,1} \cap T_2$, and identify one

circle in $\Sigma_{1,1} \cap T'_1$ with the circle in $\Sigma_{2,1} \cap T'_2$. Then we do the other identifications arbitrarily. Note that there are many ways to do the further identification, since we can isotopy one Σ_{2,k_0} such that its intersection with T_2 slides over the other circles of $\Sigma_{2,k} \cap T_2$, while all the other surfaces in $\{\Sigma_{2,k}\}$ are fixed.

It is easy to see that $i : \Sigma \looparrowright N$ is a properly immersed surface, and it satisfies conditions (1)-(5) in the proposition, by the construction.

Now we show that i is π_1 -injective. Suppose there is a map $j : S^1 \rightarrow \Sigma$ which is not null-homotopic in Σ , but $i \circ j : S^1 \rightarrow N$ is null-homotopic in N .

We can assume that $i \circ j$ is transverse to the decomposition tori $T \cup T'$, and this j minimizes the number of points in $(i \circ j)^{-1}(T \cup T') \subset S^1$. This number is not zero, otherwise, it contradicts with the π_1 -injectivity of fibered surfaces.

Since $i \circ j$ is null-homotopic, it can be extended to a map $k : D^2 \rightarrow N$ such that $k|_{S^1} = i \circ j$. We can assume that k is transverse to $T \cup T'$, and $k^{-1}(T \cup T')$ consists of simple arcs in D^2 , since we can homotopy k relative to S^1 such that there are no circle components in $k^{-1}(T \cup T')$.

Then there exists an arc $\alpha \subset S^1$ and an arc component β in $k^{-1}(T \cup T') \subset D^2$, such that α and β share end points and there are not other components of $k^{-1}(T \cup T')$ lying in the subdisc $B \subset D^2$ bounded by $\alpha \cup \beta$. Suppose that $j(\alpha)$ lies in $\Sigma_{i,k} \subset N_i$, then k also maps B into N_i . We can suppose that T is the decomposition torus containing the k -image of β . Then it is easy to see that the k -images of two end points of α lie in the same component of $\Sigma_{i,k} \cap T$, by considering the algebraic intersection number between $\Sigma_{i,k}$ and $\alpha \cup \beta$. Moreover, $k| : \beta \rightarrow T$ is homotopic to a map with image in $\Sigma_{i,k} \cap T$, relative to the boundary of β .

Then it is routine to check that $j| : \alpha \rightarrow \Sigma_{i,k}$ is homotopy to a map with image in $i^{-1}(T)$, relative to the boundary. After a further homotopy of j at a neighborhood of α , we get $j' : S^1 \rightarrow \Sigma$ which is homotopy to j and has fewer number of points in $(i \circ j')^{-1}(T \cup T') \subset S^1$.

So we get a contradiction, and $i : \Sigma \looparrowright N$ is π_1 -injective. $\square$

The following proposition proves the nonseparability of $i_*(\pi_1(\Sigma)) < \pi_1(N)$ constructed in Proposition 3.6. The proof is essentially a proof of the aspirality character defined in [Li1] being not ± 1 , but we do not use the terminology of aspirality character, since the picture is relatively simple.

Proposition 3.7. *For the properly immersed surface $i : \Sigma \looparrowright N$ constructed in Proposition 3.6, $i_*(\pi_1(\Sigma)) < \pi_1(N)$ is a nonseparable subgroup.*

Proof. Suppose that $i_*(\pi_1(\Sigma)) < \pi_1(N)$ is separable, we want to get a contradiction.

Let $\tilde{N}$ be the covering space of N corresponding to $i_*(\pi_1(\Sigma))$. Since each component of $\Sigma \cap i^{-1}(N_k)$ is a fibered surface in N_k , it is easy to see that $\tilde{N}$ is homeomorphic to $\Sigma \times \mathbb{R}$. So $i : \Sigma \looparrowright N$ lifts to an embedding $\Sigma \hookrightarrow \tilde{N}$.

Since $i_*(\pi_1(\Sigma)) < \pi_1(N)$ is separable, by [Sc], there exists an intermediate finite cover $\hat{N} \rightarrow N$ such that $i : \Sigma \looparrowright N$ lifts to an embedding $\hat{i} : \Sigma \hookrightarrow \hat{N}$.

Since $i : \Sigma \looparrowright N$ is a proper immersion, $\hat{i} : \Sigma \hookrightarrow \hat{N}$ is a proper embedding. So Σ defines a nontrivial cohomology class $\sigma \in H^1(\hat{N}; \mathbb{Z})$, by duality.

For each decomposition torus $\hat{T}_s$ in $\hat{N}$, suppose $\Sigma \cap \hat{T}_s$ consists of k_s parallel circles. Let K be the least common multiple of all k_s . Take the K -sheet cyclic cover of $\hat{N}$ along Σ (corresponding to the kernel of $H_1(\hat{N}; \mathbb{Z}) \xrightarrow{\sigma} \mathbb{Z} \rightarrow \mathbb{Z}_K$), we

get a finite cover $\bar{N} \rightarrow N$. Then Σ embeds into $\bar{N}$, and it intersects with each decomposition torus of $\bar{N}$ exactly once.

Let $\bar{N}_1$ and $\bar{N}_2$ be the geometric pieces of $\bar{N}$ containing $\Sigma_{1,1}$ and $\Sigma_{2,1}$ respectively. Since $\Sigma_{1,1} \cap \Sigma_{2,1}$ along $s \cup s'$, let $\bar{T}$ and $\bar{T}'$ be the decomposition tori in $\bar{N}_1 \cap \bar{N}_2$ corresponding to these two circles, respectively. Then the finite cover $\bar{N} \rightarrow N$ induces finite covers:

$$\begin{aligned} \bar{N}_1 &\rightarrow N_1, & \bar{N}_2 &\rightarrow N_2, \\ \bar{T} &\rightarrow T, & \bar{T}' &\rightarrow T'. \end{aligned}$$

Since $\bar{T} \rightarrow T$ and $\bar{T}' \rightarrow T'$ are both induced by $\bar{N}_1 \rightarrow N_1$ and $\bar{N}_2 \rightarrow N_2$, we will get two relations between $\deg(\bar{T} \rightarrow T)$ and $\deg(\bar{T}' \rightarrow T')$, then get a contradiction.

Since $\Sigma_{1,1}$ is an embedded fibered surface in both $\bar{N}_1$ and N_1 , $\bar{N}_1$ is a cyclic cover of N_1 along $\Sigma_{1,1}$. Similarly, $\bar{N}_2$ is a cyclic cover of N_2 along $\Sigma_{2,1}$.

Since $\Sigma_{1,1} \cap T$ consists of A parallel circles and $\Sigma_{1,1} \cap T'$ consists of B parallel circles, while $\Sigma_{1,1} \cap \bar{T}$ and $\Sigma_{1,1} \cap \bar{T}'$ are both only one circle, $\bar{N}_1 \rightarrow N_1$ is a cyclic cover with degree being a multiple of $\text{lcm}(A, B)$, and

$$(1) \quad B \cdot \deg(\bar{T} \rightarrow T) = \deg(\bar{N}_1 \rightarrow N_1) = A \cdot \deg(\bar{T}' \rightarrow T').$$

We also have that $\Sigma_{2,1}$ is an embedded fibered surface in both $\bar{N}_2$ and N_2 . Since $\Sigma_{2,1} \cap T$, $\Sigma_{2,1} \cap T'$, $\Sigma_{2,1} \cap \bar{T}$ and $\Sigma_{2,1} \cap \bar{T}'$ are all just one circle, and $\bar{N}_2 \rightarrow N_2$ is a cyclic cover, we have

$$(2) \quad \deg(\bar{T} \rightarrow T) = \deg(\bar{N}_2 \rightarrow N_2) = \deg(\bar{T}' \rightarrow T').$$

Equations (1) and (2) imply that $A = B$, which contradicts with condition (5) in Proposition 3.6. So $i_*(\pi_1(\Sigma))$ is a nonseparable subgroup of $\pi_1(N)$. $\square$

Remark 3.8. From the proof of Proposition 3.7, the readers can see that the main ingredient for the nonseparability of $\pi_1(\Sigma)$ is the subsurface $\Sigma_{1,1} \cup_{s \cup s'} \Sigma_{2,1}$. However, the author can not prove that $\pi_1(\Sigma_{1,1} \cup_{s \cup s'} \Sigma_{2,1})$ is nonseparable in $\pi_1(N)$, although it seems quite plausible.

In the proof of Proposition 3.7, we do need the properness of the immersed surface $i : \Sigma \looparrowright N$, so that we can do cyclic covering of $\hat{N}$ along Σ to get $\bar{N}$, and then get the contradiction. Actually, most part of the proof can be translated to purely group theoretical language, except that the author does not know how to interpret "properly immersed surfaces" algebraically.

3.4. Proof of Theorem 1.3.

Now we are ready to prove Theorem 1.3.

Proof. Suppose that M supports one of eight Thurston's geometries. If M supports the S^3 - or $S^2 \times \mathbb{E}^1$ -geometry, LERFness trivially holds. If M supports the $\mathbb{E}^3$ -, Nil-, $\mathbb{H}^2 \times \mathbb{E}^1$ - or $\text{PSL}_2(\mathbb{R})$ -geometry, LERFness is proved in [Sc]. If M supports the Sol-geometry, a proof of LERFness can be found in [NW2]. If M is a hyperbolic 3-manifold, LERFness is shown by the celebrated works of Wise ([Wi] for cusped case) and Agol ([Ag2] for closed case).

Now we need to show that non-geometric 3-manifolds have nonLERF fundamental groups. We first suppose that M is a mixed 3-manifold, i.e. there is a hyperbolic piece in M .

If M is not a closed manifold, then Lemma 3.4 and Lemma 3.5 imply that M has a finite semicover $N = N_1 \cup_{T \cup T'} N_2$ satisfying the conditions in Lemma 3.5. In particular, $\pi_1(N)$ is a subgroup of $\pi_1(M)$. Then Proposition 3.6 constructs a

non-closed surface subgroup (free subgroup) $\pi_1(\Sigma)$ in $\pi_1(N)$, and Proposition 3.7 shows that $\pi_1(\Sigma)$ is not separable in $\pi_1(N)$. Finally, Lemma 2.3 implies that $\pi_1(\Sigma)$ is not separable in $\pi_1(M)$, thus $\pi_1(M)$ is not LERF.

If M is a closed mixed 3-manifold, then the above proof also shows that there is a nonseparable free subgroup in $\pi_1(M)$. We need also to construct a nonseparable closed surface subgroup.

Let $N \rightarrow M$ be the finite semicover constructed in Lemma 3.5 (with $\partial N \neq \emptyset$), and $\Sigma \looparrowright N$ be the π_1 -injective properly immersed surface constructed in Proposition 3.6. To make the geometric picture simpler, we use Lemma 3.2 to find a finite cover M' of M , such that N lifts to an embedded submanifold of M' .

In this case, $i_1 : \Sigma \looparrowright M'$ is an immersed subsurface, but is not properly immersed. So we can not use the proof of Proposition 3.7 for this Σ . Now we need to extend Σ to a closed surface Σ' , with an immersion $i_2 : \Sigma' \looparrowright M'$. Then we can apply a similar argument as in the proof of Proposition 3.7.

The construction of $i_2 : \Sigma' \looparrowright M'$ is already done in Section 8 of [Li1], so we only give a sketch here.

Let the boundary components of Σ be $s_1, \dots, s_m$, with each s_i lying on a decomposition torus $T_i \subset M'$. By Theorem 4.11 of [DLW], there exists an essentially immersed surface $R_i \looparrowright M'$, such that ∂R_i consists of two components b_i and $\bar{b}_i$, with b_i and $\bar{b}_i$ mapped to a positive and a negative multiple of $s_i \subset T_i$ respectively, with the same covering degree. Moreover, a neighborhood of ∂R_i in R_i is mapped to the side of T_i other than N , and R_i intersects with $\mathcal{T}_{M'}$ minimally.

Then we can take some finite cover of Σ and take another copy with the opposite orientation. Together with proper number of copies of $R_i, i = 1, \dots, m$, they can be pasted together to get a π_1 -injective immersed closed surface $\Sigma' \looparrowright M'$. Then a similar argument as in Proposition 3.7 shows that $\pi_1(\Sigma')$ is not separable in $\pi_1(M')$, so also not separable in $\pi_1(M)$.

If M is a graph manifold, it was already showed in [NW2] that $\pi_1(M)$ is not LERF. So we do not give the detail of the construction of nonseparable surface subgroups.

The first step of the construction is to show that M has a finite semicover $N = S \times I/\phi$, where $S = S_1 \cup_{c \cup c'} S_2$ and ϕ is a composition of Dehn twists along c and c' . Then we perturb the fibered structures on both N_1 and N_2 (since Seifert fibered spaces have less flexible fibered structures) to get a π_1 -injective properly immersed subsurface similar to what we get in Proposition 3.6. Then a similar argument as in Proposition 3.7 shows that this surface subgroup is not separable. Here we need to use the fact that two adjacent Seifert pieces in a graph manifold have incompatible regular fibers on their intersection.

However, it seems not easy to construct a nonseparable closed surface subgroup in a general closed graph manifold. $\square$

Remark 3.9. In [NW1], it is shown that the construction in [RW] gives π_1 -injective properly immersed subsurface $\Sigma \looparrowright M$ in some graph manifold M , such that $\pi_1(\Sigma)$ is not contained in any finite index subgroup of $\pi_1(M)$ (not engulfed). In the proof of [NW1], it is only used the fact that the surfaces constructed in [RW] have the infinite plane property. Since the surfaces we constructed in the proof of Theorem 1.3 also have the infinite plane property, for any mixed 3-manifold, we can find a

finite cover M of it and a π_1 -injective properly immersed subsurface $\Sigma \looparrowright M$ such that $\pi_1(\Sigma)$ is not contained in any finite index subgroup of $\pi_1(M)$.

In [NW2], it is shown that all graph manifold groups contain

$$L = \langle x, y, r, s \mid x^r = x, y^r = y, x^s = x \rangle$$

as a subgroup. Then the nonLERFness of L implies the nonLERFness of graph manifold groups. It is easy to see that some mixed manifolds, e.g. double of any cusped hyperbolic 3-manifold, do not have L as a subgroup in their fundamental groups. So L is not the source of the nonLERFness of these groups.

Since free products of LERF groups are still LERF, we have the following direct corollary of Theorem 1.3.

Corollary 3.10. *Let M be a compact 3-manifold with empty or tori boundary, then $\pi_1(M)$ is LERF if and only if all the prime factors of M support one of Thurston's eight geometries.*

The knot complements in S^3 is also a classical family of interesting 3-manifolds, and each knot is either a torus knot, or a hyperbolic knot, or a satellite knot. We have the following corollary for this family.

Corollary 3.11. *Let M be the complement of a knot $K \subset S^3$, then $\pi_1(M)$ is LERF if and only if K is a torus knot or a hyperbolic knot.*

4. NONLERFNESS OF GROUPS OF HYPERBOLIC 3-MANIFOLDS UNION ALONG A CIRCLE

In this section, we will give the proof of Theorem 1.4. The proof is very similar to the proof of Theorem 1.3. For some lemmas and propositions in this section, we will only give a sketch of the proof, and point out the necessary modifications of the corresponding proofs in Section 3.

In the proof of the nonLERFness of $M_1 \cup_{S^1} M_2$, we actually only use properties of hyperbolic 3-manifolds for M_1 , and do not have much requirement for M_2 . So we will state some more general results on nonLERFness of $\mathbb{Z}$ -amalgamated groups in Subsection 4.2.

4.1. NonLERFness of $\pi_1(M_1 \cup_{S^1} M_2)$ for hyperbolic 3-manifolds M_1 and M_2 . Suppose that M_1 and M_2 are two finite volume hyperbolic 3-manifolds (possibly with cusps), and $i_k : S^1 \rightarrow M_k, k = 1, 2$ be two essential circles. Here we can assume that both i_k are embeddings in $\text{int}(M_k)$, and denote the image of i_k by γ_k . It is possible that the element in $\pi_1(M_k)$ corresponding to γ_k is a parabolic element or a nonprimitive element. However, for most of the time, the readers can think γ_k as a simple closed geodesic in M_k .

Let $X = M_1 \cup_\gamma M_2$ be the space obtained by identifying γ_1 and γ_2 by a homeomorphism, then we need to show that $\pi_1(X)$ is not separable. For a standard graph of space, the edge space should be $S^1 \times I$. Here we directly paste these two manifolds together along the circles, which makes the picture simpler. We also give orientations on γ_1 and γ_2 such that their orientations agree with each other under the identification.

For any point in X , either it has a neighborhood homeomorphic to B^3 or B_+^3 , or it has a neighborhood homeomorphic to a union of two B^3 s along $I_z = B^3 \cap (z\text{-axis})$, i.e. $B^3 \cup_{I_z} B^3$.

We first give a name for spaces locally look like B^3 , B_+^3 or $B^3 \cup_I B^3$.

Definition 4.1. A compact Hausdorff space X is called a *singular 3-manifold* if for any point $x \in X$, either it has a neighborhood homeomorphic to B^3 or B_+^3 , or it has a neighborhood homeomorphic to $B^3 \cup_{I_z} B^3$ with $x \in I_z$. We call points in the first class *regular points*, and points in the second class *singular points*.

We can think about a singular 3-manifold X as a union of finitely many 3-manifolds along disjoint simple closed curves, and we call each of these 3-manifolds a 3-manifold piece of X .

In the proof of Theorem 1.3, the concept of finite semicover played an important role, so we need to define a corresponding concept for singular 3-manifolds. Here the set of singular points in singular 3-manifolds correspond to the set of decomposition tori in 3-manifolds.

Definition 4.2. Let Y, Z be two singular 3-manifolds, a map $i : Y \rightarrow Z$ is called a *singular finite semicover* if for any point $y \in Y$, one of the following case holds.

- (1) i maps a neighborhood of y to a neighborhood of $i(y)$ by homeomorphism.
- (2) y is a regular point and $i(y)$ is a singular point, such that i maps a B^3 neighborhood of y to one of the B^3 in a $B^3 \cup_{I_z} B^3$ neighborhood of $i(y)$ by homeomorphism.

So a singular finite semicover maps singular points to singular points, and maps most of regular points to regular points. It maps each 3-manifold piece of Y to a 3-manifold piece of Z by a finite cover.

It is easy to see that a singular finite semicover $i : Y \rightarrow Z$ induces an injective homomorphism on the fundamental groups. The author also expects that a singular finite semicover gives a separable subgroup $\pi_1(Y) < \pi_1(Z)$, but we do not need this result here.

The following lemma corresponds to Lemma 3.4.

Lemma 4.3. *Let $X = M_1 \cup_\gamma M_2$ be a union of two finite volume hyperbolic 3-manifolds along an essential circle, there exists a singular 3-manifold $Y = N_1 \cup_c N_2$ such that the following hold.*

- (1) Y is a union of two hyperbolic 3-manifolds N_1 and N_2 , where N_k is a finite cover of M_k ($k = 1, 2$), and the set of singular points is one oriented circle.
- (2) Each N_k is a fibered 3-manifold with a fixed fibered surface S_k , such that the algebraic intersection number $[S_k] \cap [c] = 1$, for $k = 1, 2$.
- (3) Y is a singular finite semicover of X , so $\pi_1(Y)$ is a subgroup of $\pi_1(X)$.

Proof. By Agol's virtual fibering theorem and virtual infinite betti number theorem ([Ag2]), there exists a finite cover M'_1 of M_1 such that M'_1 is a fibered 3-manifold and $b_1(M'_1) > 1$. Let $\gamma'_1 \subset M'_1$ be one oriented elevation (component of preimage) of $\gamma_1 \subset M_1$. If γ'_1 is nullhomologous in M'_1 , we can use Theorem 2.5 to find a further finite cover M''_1 such that γ'_1 lifts to a non null-homologous curve in M''_1 .

Since the fibered cone is an open set in $H^1(M''_1; \mathbb{R})$, there exists a fibered surface S_1 in M''_1 which has positive intersection number with γ'_1 . So we have $[S_1] \cap [\gamma'_1] = a_1 \in \mathbb{Z}_+$, and $\deg(\gamma'_1 \rightarrow \gamma_1) = b_1$.

By the same construction, we get a finite cover $M''_2 \rightarrow M_2$ with a fibered surface S_2 , such that for some oriented elevation γ'_2 of γ_2 , $[S_2] \cap [\gamma'_2] = a_2 \in \mathbb{Z}_+$ and $\deg(\gamma'_2 \rightarrow \gamma_2) = b_2$.

Let N_1 be the $a_1 b_2$ -sheet cyclic cover of M''_1 along S_1 , and c_1 be one elevation of γ'_1 . Then $[S_1] \cap [c_1] = 1$ and $\deg(c_1 \rightarrow \gamma_1) = b_1 b_2$. Similarly, let N_2 be the $a_2 b_1$ -sheet

cyclic cover of M_2'' along S_2 , and c_2 be one elevation of γ_1' . Then $[S_2] \cap [c_2] = 1$ and $\deg(c_2 \rightarrow \gamma_2) = b_1 b_2$.

Since $c_1 \rightarrow \gamma_1$ and $c_2 \rightarrow \gamma_2$ have the same degree, we can identify c_1 and c_2 (as oriented curves) to get the desired singular finite semicover $Y = N_1 \cup_c N_2$. $\square$

Remark 4.4. Actually, we may get a result as strong as in Lemma 3.4, i.e. $Y = N_1 \cup_c N_2$ is an $S_1 \vee S_2$ bundle over S^1 . However, we did not state this statement in the lemma. One reason is that we need to homotopy the curve c_k in N_k to get this fibered structure. Moreover, each closed curve c_k may not be a (virtually) closed orbit of the pseudo-Anosov suspension flow of a (virtual) fibered structure of N_k . So it is not a natural object, from the dynamical point of view. Of course, stating this result in the form of Lemma 4.3 does not affect the proof of Theorem 1.4.

For simplicity, we still use $X = M_1 \cup_\gamma M_2$ to denote the singular 3-manifold obtained in Lemma 4.3. Then we have the following lemma corresponding to Lemma 3.5.

Lemma 4.5. *For the singular 3-manifold $X = M_1 \cup_\gamma M_2$ constructed in Lemma 4.3, there exists a singular 3-manifold $Y = N_1 \cup_{c \cup c'} N_2$ such that the following hold.*

- (1) *Y is a union of two hyperbolic 3-manifolds N_1 and N_2 , where each N_k is a finite cover of M_k ($k = 1, 2$), and the set of singular points is a union of two oriented circles.*
- (2) *The homomorphism $H_1(c \cup c'; \mathbb{Z}) \rightarrow H_1(N_1; \mathbb{Z})$ induced by inclusion is injective.*
- (3) *For each $N_k, k = 1, 2$, there exists a fibered surface $S'_k \subset N_k$ such that the algebraic intersection number $[S'_k] \cap [c] = [S'_k] \cap [c'] = 1$.*
- (4) *Y is a finite semicover of X , so $\pi_1(Y)$ is a subgroup of $\pi_1(X)$.*

Proof. Let γ_i be the oriented copy of γ in M_i .

By a similar argument as in the proof of Lemma 3.5, and using the virtual retract property of a $\mathbb{Z} * \mathbb{Z} = \langle \pi_1(\gamma), g^n \pi_1(\gamma) g^{-n} \rangle$ subgroup in $\pi_1(M_1)$, we can find a finite cover N_1 of M_1 and two distinct homeomorphic liftings c_1 and c'_1 of $\gamma_1 \subset M_1$, such that $H_1(c_1 \cup c'_1; \mathbb{Z}) \rightarrow H_1(N_1; \mathbb{Z})$ is injective.

In the conclusion of Lemma 4.3, we fixed a fibered surface S_1 in M_1 whose algebraic intersection number with γ_1 is 1. For an elevated fibered surface $S'_1 \subset N_1$, the algebraic intersection numbers of S'_1 with both c_1 and c'_1 are both equal to 1.

By doing a similar construction for M_2 (actually a simpler construction works since we do not require condition (2) for N_2), we get a finite cover N_2 of M_2 , with two homeomorphic liftings c_2 and c'_2 of γ_2 , and a fibered surface S'_2 of N_2 with $[S'_2] \cap [c_2] = [S'_2] \cap [c'_2] = 1$.

Then we paste N_1 and N_2 together by identifying c_1 with c_2 (denoted by c) and identifying c'_1 and c'_2 (denoted by c') to get the desired singular semicover Y . $\square$

For singular 3-manifolds, we need a definition in this singular world that corresponds to immersed surfaces in 3-manifolds.

We first define singular surfaces, which plays the same role for surfaces in 3-manifolds. This definition is very similar to the definition of singular 3-manifolds.

Definition 4.6. A compact Hausdorff space K is called a *singular surface* if for any point $k \in K$, either it has a neighborhood homeomorphic to B^2 or B^2_+ , or it has a neighborhood homeomorphic to $B^2 \vee B^2$, with k lying in the intersection of

two discs. We call the points in the first class *regular points*, and points in the second class *singular points*.

We can think a singular surface K as a union of finitely many compact surfaces, pasting along finitely many points in the interior. We call each of these surfaces a surface piece of K .

Then we can define singular immersions from singular surfaces to singular 3-manifolds.

Definition 4.7. Let $i : K \rightarrow X$ be a map from a singular surface to a singular 3-manifold. We say that i is a *singular immersion* if the following conditions hold.

- (1) i maps the singular set of K to the singular set of X .
- (2) The restriction of i on each surface piece of K is a proper immersion from the surface to a 3-manifold piece of X .
- (3) For any singular point $k \in K$, there exist a $B^2 \vee B^2$ neighborhood of k and a $B^3 \cup_{I_\pm} B^3$ neighborhood of $i(k)$, such that i maps two B^2 s to distinct B^3 s in $B^3 \cup_{I_\pm} B^3$, and each B^2 is mapped to the intersection of B^3 with the xy -plane by homeomorphism.

Now we state the corresponding result of Proposition 3.6. Here we do not define the notion of "singular proper immersion", since it is quite subtle in the current context.

Proposition 4.8. *For the singular 3-manifold $Y = N_1 \cup_{c \cup c'} N_2$ and fibered surfaces $S'_k \subset N_k$ constructed in Lemma 4.5, there exists a singular surface K and a π_1 -injective singular immersion $i : K \rightarrow Y$ such that the following hold.*

- (1) K is a union of oriented connected subsurfaces as $K = (\Sigma_{1,1} \cup \Sigma_{1,2}) \cup (\cup_{k=1}^{2n} \Sigma_{2,k})$, with $i(\Sigma_{1,j}) \subset N_1$ and $i(\Sigma_{2,k}) \subset N_2$.
- (2) There are $4n$ singular points in K , each singular point lies in some $\Sigma_{1,j} \cap \Sigma_{2,k}$, and each $\Sigma_{2,k}$ contains exactly two singular points.
- (3) The restriction of i on $\Sigma_{1,j}$ s and $\Sigma_{2,k}$ s are all embeddings, and the images are fibered surfaces of N_1 and N_2 respectively.
- (4) Each $\Sigma_{2,k}$ is a copy of S'_2 in N_2 , and the two singular points in $\Sigma_{2,k}$ are mapped to the intersection of $\Sigma_{2,k}$ with c and c' respectively.
- (5) $\Sigma_{1,1} \cap \Sigma_{2,1}$ consists of two singular points p and p' , with $i(p) \in c$ and $i(p') \in c'$.
- (6) The algebraic intersection number $[\Sigma_{1,1}] \cap [c_1] = A$ and $[\Sigma_{1,1}] \cap [c'_1] = B$, with $A \neq B$; while the algebraic intersection number $[\Sigma_{1,2}] \cap [c_2] = 2n - A$ and $[\Sigma_{1,2}] \cap [c'_2] = 2n - B$. Here c_i and c'_i are the oriented copies of c and c' in N_i respectively, and all these algebraic intersection numbers are positive integers.
- (7) The set $\{\text{singular points in } \Sigma_{1,1}\} \cap i^{-1}(c)$ has cardinality A . Suppose this set is $\{a_1, \dots, a_A\}$, then $\Sigma_{1,1}$ has positive local intersection number with c_1 at each a_l . The same statement holds for $\{\text{singular points in } \Sigma_{1,1}\} \cap i^{-1}(c')$ (with A replaced by B), $\{\text{singular points in } \Sigma_{1,2}\} \cap i^{-1}(c)$ and $\{\text{singular points in } \Sigma_{1,2}\} \cap i^{-1}(c')$ (with A replaced by $2n - A$ and $2n - B$, respectively).
- (8) For each $l \in \{1, \dots, A\}$, take the embedded oriented subarc of c from $i(a_1)$ to $i(a_l)$. Then slightly move it along the positive direction of c_1 , to get an oriented arc ρ_l with end points away from $\Sigma_{1,1}$. Then the algebraic

intersection number of $\Sigma_{1,1}$ with ρ_l is equal to $l - 1$. Similar statements also hold for $\{\text{singular points in } \Sigma_{1,1}\} \cap i^{-1}(c')$, $\{\text{singular points in } \Sigma_{1,2}\} \cap i^{-1}(c)$ and $\{\text{singular points in } \Sigma_{1,2}\} \cap i^{-1}(c')$.

This proposition looks more complicated than Proposition 3.6, and we give some remarks here.

Remark 4.9. The conditions (1)-(6) in Proposition 4.8 correspond to the conditions in Proposition 3.6, and conditions (7) and (8) in Proposition 4.8 correspond to the "properness" of this singular immersion. Although we do not have the condition

$$i^{-1}\{\text{singular points in } Y\} = \{\text{singular points in } K\},$$

conditions (6) and (7) imply that the total intersection number of $\Sigma_{1,j}$ with c_1 at the points in $(i^{-1}(c) \setminus \{\text{singular points in } \Sigma_{1,j}\}) \cap \Sigma_{1,j}$ is zero, and it also holds for c' . So it is a weak and algebraic version of $i^{-1}(c \cup c') \cap \Sigma_{1,j} = \{\text{singular points in } \Sigma_{1,j}\}$.

Here we use algebraic intersection number instead of geometric intersection number (or number of components in the intersection) as in Proposition 3.6. For a fibered surface and a closed orbit of the pseudo-Anosov suspension flow (or a boundary component of the 3-manifold), the algebraic intersection number is always equal to the geometric intersection number (or number of components in the intersection). However, the circles c_1 and c'_1 in N_1 may not be (virtually) closed orbits of the pseudo-Anosov suspension flow, even up to homotopy. Although we can homotopy c_1 and c'_1 such that their algebraic intersection numbers with one fibered surface are equal to geometric intersection numbers, but there are two fibered surfaces $\Sigma_{1,1}$ and $\Sigma_{1,2}$ in N_1 , and we may not be able to do it simultaneously for both $\Sigma_{1,1}$ and $\Sigma_{1,2}$.

Proof. By the same argument as in the proof of Proposition 3.6, we can construct two fibered surfaces $\Sigma_{1,1}$ and $\Sigma_{1,2}$ in N_1 such that condition (6) holds. Take $2n$ copies of S'_2 in N_2 , and denote them by $\Sigma_{2,1}, \dots, \Sigma_{2,2n}$.

First suppose we choose any $A, B, 2n - A, 2n - B$ points in $\Sigma_{1,1} \cap c_1, \Sigma_{1,1} \cap c'_1, \Sigma_{1,2} \cap c_1, \Sigma_{1,2} \cap c'_1$ respectively, such that the corresponding surfaces and curves have positive local intersection numbers at these points. If we identify these points with $(\cup_{k=1}^{2n} \Sigma_{2,k}) \cap (c_2 \cup c'_2)$ by an arbitrary way, we get a singular surface K and a singular immersion satisfying conditions (1)-(4), (6) and (7).

So we need to choose these points carefully so that condition (8) holds, and then do the correct pasting such that condition (5) holds.

The choice of these four families of points follows the same process, so we only consider $\Sigma_{1,1} \cap c_1$. Although the algebraic intersection number between $\Sigma_{1,1}$ and c_1 is A , there might be more geometric intersection points. So we assume that there are $A + 2m$ intersection points in $\Sigma_{1,1} \cap c_1$. Take any positive intersection point a'_1 in $\Sigma_{1,1} \cap c_1$. By following the orientation of c_1 , we denote the other points of $\Sigma_{1,1} \cap c_1$ by $a'_2, \dots, a'_{A+2m}$. For any $l \in \{1, \dots, A+2m\}$, take the embedded oriented subarc in c_1 from a'_1 to a'_l , then move it slightly along the positive direction of c_1 , and denote it by ρ'_l . Whenever we move from a'_l to a'_{l+1} , the algebraic intersection number $[\Sigma_{1,1}] \cap [\rho'_l]$ differs from $[\Sigma_{1,1}] \cap [\rho'_{l+1}]$ by 1 or -1 , depending on whether $\Sigma_{1,1}$ intersects with c_1 positively or negatively at a'_{l+1} . Since $[\Sigma_{1,1}] \cap [\rho'_1] = 0$ and $[\Sigma_{1,1}] \cap [\rho'_{A+2m}] = A - 1$, it is easy to find A points in $\{a'_1, a'_2, \dots, a'_{A+2m}\}$ (with $a_1 = a'_1$), such that they are all positive intersection points and satisfy condition (8).

Then we can paste the $2n$ points in $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap c_1$ (and $(\Sigma_{1,1} \cup \Sigma_{1,2}) \cap c'_1$) chosen above with the $2n$ points in $(\cup_{k=1}^{2n} \Sigma_{2,k}) \cap c_2$ (and $(\cup_{k=1}^{2n} \Sigma_{2,k}) \cap c'_2$), to get a singular immersed surface $i : K \looparrowright Y$. By doing isotopy of $\Sigma_{2,1}$ in N_2 , we can make sure the pasting satisfies condition (5).

The π_1 -injectivity of $i : K \looparrowright Y$ follows from the same π_1 -injectivity argument in Lemma 3.6. Note that we do need condition (8) here. $\square$

Then we can show that the above π_1 -injective singular immersion gives a non-separable subgroup in $\pi_1(Y)$. The proof is similar to the proof of Proposition 3.7.

Proposition 4.10. *For the singular immersion $i : K \looparrowright Y$ constructed in Proposition 4.8, $i_*(\pi_1(K)) < \pi_1(Y)$ is a nonseparable subgroup.*

Proof. Suppose that $i_*(\pi_1(K)) < \pi_1(Y)$ is separable, then we want to get a contradiction.

Since each surface piece of K is mapped to a fibered surface in the corresponding 3-manifold piece of Y , the covering space $\tilde{Y}$ of Y corresponding to $\pi_1(K)$ is homeomorphic to a union of $\Sigma_{1,j} \times \mathbb{R}$ (with $j = 1, 2$) and $\Sigma_{2,k} \times \mathbb{R}$ (with $k = 1, \dots, 2n$), by pasting along preimages of c_i and c'_i (with $i = 1, 2$). In particular, $i : K \looparrowright Y$ lifts to an embedding in $\tilde{Y}$. By the separability of $i_*(\pi_1(K))$ and [Sc], there exists an intermediate finite cover $p : \hat{Y} \rightarrow Y$ of $\tilde{Y}$ such that $i : K \looparrowright Y$ lifts to an embedding $\hat{i} : K \hookrightarrow \hat{Y}$.

In Proposition 3.7, we took a finite cyclic cover of $\hat{N}$ along Σ . It can be done either geometrically, i.e. take finitely many copies of $\hat{N} \setminus \Sigma$ and paste together, or algebraically, i.e. take a finite cyclic cover dual to the cohomology class defined by Σ .

Here we do not follow the geometric process, since $\hat{i}^{-1}\{\text{singular points in } \hat{Y}\} \neq \{\text{singular points in } K\}$, although we can make it holds by homotopy the attaching circles $p^{-1}(c \cup c')$ in 3-manifold pieces of $\hat{Y}$.

Now we show that $K \subset \hat{Y}$ defines a cohomology class $\kappa \in H^1(Y; \mathbb{Z})$, by using duality, i.e. taking algebraic intersection number.

Since K intersects with each 3-manifold piece in $\tilde{Y}$, it also intersects with each 3-manifold piece in $\hat{Y}$. For each 3-manifold piece $\hat{N}_s$ of $\hat{Y}$, $K \cap \hat{N}_s$ is a properly embedded oriented surface in $\hat{N}_s$, so it defines a cohomology class $\kappa_s \in H^1(\hat{N}_s; \mathbb{Z})$.

For each component $\hat{c}$ of $p^{-1}(c \cup c')$, suppose that it is adjacent to $\hat{N}_1$ and $\hat{N}_2$, then we need to show that $\kappa_1|_{\hat{c}} = \kappa_2|_{\hat{c}}$, i.e. the algebraic intersection numbers $[\hat{N}_1 \cap K] \cap [\hat{c}]$ and $[\hat{N}_2 \cap K] \cap [\hat{c}]$ are equal to each other. We can suppose that $\hat{N}_1$ and $\hat{N}_2$ cover N_1 and N_2 respectively.

Since $\Sigma_{1,1}$ and $\Sigma_{1,2}$ are different fibered surfaces of N_1 , only one of them lies in $\hat{N}_1$. Without loss of generality, we suppose that $K \cap \hat{N}_1 = \Sigma_{1,1}$, and $\hat{c}$ is a component of $p^{-1}(c)$. All the other cases follow from the same argument.

Since $\Sigma_{1,1}$ is a fibered surface in both $\hat{N}_1$ and N_1 , $\hat{N}_1 \rightarrow N_1$ is a finite cyclic cover dual to $\Sigma_{1,1}$, and let the covering degree be D . Recall that $[\Sigma_{1,1}] \cap [c_1] = A$. Then $p^{-1}(c) \cap \hat{N}_1$ has $\gcd(A, D)$ many components ($\hat{c}$ is one of them), and each of them has algebraic intersection number $\frac{A}{\gcd(A, D)}$ with $\Sigma_{1,1}$.

So we have shown that $\langle \kappa_1, \hat{c} \rangle = \frac{A}{\gcd(A, D)}$, and need to show $\langle \kappa_2, \hat{c} \rangle = \frac{A}{\gcd(A, D)}$.

We first show that for $\hat{i} : \Sigma_{1,1} \rightarrow \hat{N}_1$, there are exactly $\frac{A}{\gcd(A,D)}$ points in $\{a_1, \dots, a_A\}$ mapped to $\hat{c}$. For two points $a_s, a_t \in \{a_1, \dots, a_A\}$, if $\hat{i}(a_s)$ and $\hat{i}(a_t)$ lie in the same component of $p^{-1}(c) \cap \hat{N}_1$, there is an oriented subarc τ of $p^{-1}(c)$ from $\hat{i}(a_s)$ to $\hat{i}(a_t)$. Take an oriented path γ in $\Sigma_{1,1}$ from a_s to a_t . Then by projecting the loop $\tau \cdot \hat{i}(\gamma^{-1})$, we get a loop δ in N_1 .

Since $\tau \cdot \hat{i}(\gamma^{-1})$ is a loop in $\hat{N}_1$, the algebraic intersection number of $\Sigma_{1,1}$ with δ is a multiple of D . On the other hand, δ consists of the projection of τ and $\hat{i}(\gamma^{-1})$ in N_1 . This enables us to compute the algebraic intersection number by another way. Since γ lies in $\Sigma_{1,1}$, its projection in N_1 has 0 algebraic intersection number with $\Sigma_{1,1}$. Since $p \circ \tau$ is a path on c_1 with initial point a_s and terminal point a_t , by condition (8) in Proposition 4.8, the algebraic intersection number of $\Sigma_{1,1}$ with $p \circ \tau$ is $nA + (t - s)$ for some $n \in \mathbb{Z}$. (In this paragraph, we may need to slightly move τ and $\hat{i}(\gamma^{-1})$ along the positive direction of the corresponding component of $p^{-1}(c)$, such that their endpoints are away from $\Sigma_{1,1}$.)

From the two ways of computing the algebraic intersection number between $\Sigma_{1,1}$ and δ , we get that $mD = nA + (t - s)$ holds for some integers m and n . So $t - s$ is a multiple of $\gcd(A, D)$. It implies that for each component of $p^{-1}(c) \cap \hat{N}_1$, there are exactly $\frac{A}{\gcd(A,D)}$ points in $\{a_1, \dots, a_A\}$ mapped to it. In particular, it holds for $\hat{c}$.

Since each of the $\frac{A}{\gcd(A,D)}$ points in $\{a_1, \dots, a_A\} \cap \hat{c}$ lies in a fibered surface in $K \cap \hat{N}_2$, and each of them has algebraic intersection number 1 with $\hat{c}$, so $\langle \kappa_2, \hat{c} \rangle = \frac{A}{\gcd(A,D)} = \langle \kappa_1, \hat{c} \rangle$.

By an M-V sequence argument, we get that $\hat{i} : K \hookrightarrow \hat{Y}$ defines a cohomology class $\kappa \in H^1(Y; \mathbb{Z})$, by taking the algebraic intersection number of any 1-cycle in Y with K .

Then we take a finite cover of $\hat{Y}$ dual to κ , as in Proposition 3.7, to get a further finite cover $q : \bar{Y} \rightarrow Y$ such that K embeds in $\bar{Y}$, and each component of $q^{-1}(c \cup c')$ intersects with exactly two surface pieces in K , with algebraic intersection number 1. Then the same argument as in the proof of Proposition 3.7 gives a contradiction, which computes the covering degrees of two components in $q^{-1}(c \cup c')$ corresponding to $\Sigma_{1,1} \cap \Sigma_{2,1}$ by two different ways. $\square$

Now we are ready to prove Theorem 1.4.

Proof. For a singular 3-manifold $X = M_1 \cup_\gamma M_2$, Lemma 4.3 and Lemma 4.5 imply that there exists a singular finite semicover $Y = N_1 \cup_{c \cup c'} N_2$ of X such that the conditions in Lemma 4.5 hold.

By Proposition 4.8, there exists a π_1 -injective singular immersion $i : K \looparrowright Y$ satisfying the conditions in Proposition 4.8. Then Proposition 4.10 implies that $i_*(\pi_1(K))$ is not separable in $\pi_1(Y)$. Since $\pi_1(Y)$ is a subgroup of $\pi_1(X)$, Lemma 2.3 implies that $i_*(\pi_1(K))$ is not separable in $\pi_1(X)$.

If both M_1 and M_2 are bounded 3-manifolds, K is a union of bounded surfaces along finitely many points, so $\pi_1(K)$ is a free group. If at least one of M_1 and M_2 is a closed 3-manifold, then K is a union of closed surfaces and (possibly empty set of) bounded surfaces along finitely many points, so $\pi_1(K)$ is a free product of free groups and closed surface groups. $\square$

The following direct corollary of Theorem 1.4 implies that the HNN extension of a hyperbolic 3-manifold group along cyclic subgroups is not LERF.

The readers may compare this corollary with the result in [Ni2], which gives a sufficient and necessary condition for an HNN extension of a free group along cyclic subgroups being LERF. Note that Niblo's condition holds for a generic pair of cyclic subgroups in a free group.

Corollary 4.11. *Let M be a finite volume hyperbolic 3-manifold, and $A, B < \pi_1(M)$ be two infinite cyclic subgroups with an isomorphism $\phi : A \rightarrow B$, then the HNN extension*

$$\pi_1(M) *_{A^t=B} = \langle \pi_1(M), t \mid tat^{-1} = \phi(a), \forall a \in A \rangle$$

is not LERF.

Proof. Let $\pi_1(M)_{A^t=B} \rightarrow \mathbb{Z}_2$ be the homomorphism which kills all elements in $\pi_1(M)$ and maps t to $\bar{1} \in \mathbb{Z}_2$. Then the kernel H is an index two subgroup of $\pi_1(M) *_{A^t=B}$.

Then H has a graph of group structure such that the graph has two vertices, and two edges connecting these two vertices. The vertex groups are two copies of $\pi_1(M)$, and the edge groups are both infinite cyclic. So H contains a subgroup which is a $\mathbb{Z}$ -amalgamation of two copies of $\pi_1(M)$. Then Theorem 1.4 implies that H is not LERF, and Lemma 2.3 implies $\pi_1(M)_{A^t=B}$ is not LERF. $\square$

4.2. More general cases. Actually, the proof of Theorem 1.4 only uses the power of hyperbolic 3-manifolds for M_1 , and M_2 only need to satisfy some mild conditions. So we have the following generalization of Theorem 1.4.

Theorem 4.12. *Let M_1 be a finite volume hyperbolic 3-manifold, and M_2 be a compact fibered manifold over the circle, i.e. $M_2 = N \times I / \phi$ for some orientation preserving self-homeomorphism $\phi : N \rightarrow N$ on a compact n -manifold N ($n > 0$). We also suppose that $\pi_1(N)$ has some nontrivial finite quotient.*

Let $S^1 \rightarrow M_1$ be an essential circle in M_1 , and $S^1 \rightarrow M_2$ be an essential circle which has nonzero algebraic intersection number with N . Then the $\mathbb{Z}$ -amalgamation

$$\pi_1(M_1 \cup_{S^1} M_2)$$

is not LERF.

We give a sketchy proof parallel to the proof of Theorem 1.4.

Proof. At first, we can find a singular finite semicover $M'_1 \cup_\gamma M'_2$ such that similar conditions in Lemma 4.3 hold. For M_1 , we still use the virtual fibered theorem, virtual infinite betti number theorem, and the virtual retract property to find a fibered structure in some finite cover of M_1 , such that conditions (1) and (2) in Lemma 4.3 hold. For M_2 , it already has a fibered structure, and we may only need to take a cyclic finite cover of M_2 along N .

Then we can finite a further singular semicover $N_1 \cup_{c \cup c'} N_2$ such that similar conditions in Lemma 4.5 hold. For M'_1 , we still use the virtual retract property and LERFness to get a finite cover satisfying conditions (2) and (3) in Lemma 4.5. For M'_2 , we use the fact that $\pi_1(N)$ admits a nontrivial finite quotient to find a finite cover N_2 of M'_2 , such that the preimage of the γ in M'_2 contains at least two components, and they have the same algebraic intersection number with an

elevation of N . We may also need to take a finite cyclic cover of N_1 along the fibered surface.

The construction of the π_1 -injective immersed singular object (not a singular surface if $n \neq 2$) in Proposition 4.8 only perturbs fibered structures in N_1 , do all nontrivial works over there, and always use the original fibered structure of N_2 . So the same construction gives a π_1 -injective immersed singular object in $N_1 \cup_{c \cup c'} N_2$, which satisfies the conditions in Proposition 4.8.

The proof of Proposition 4.10 does not use any 3-manifold topology. It only uses the fiber bundle over circle structures and elementary counting of covering degrees. So the same proof shows that the above π_1 -injective immersed singular object gives nonseparable subgroup in $\pi_1(N_1 \cup_{c \cup c'} N_2)$, and also in $\pi_1(M_1 \cup_{S^1} M_2)$.

The nonseparable subgroup constructed above is a free product of surface groups, finite index subgroups of $\pi_1(N)$ and free groups. $\square$

Since the perturbation of fiber bundle over circle structures works in any dimension, we have the following corollary. The conditions in this corollary is easier to check.

Corollary 4.13. *Let M_1 be a finite volume hyperbolic 3-manifold, and M_2 be a compact manifold with a fiber bundle over circle structure and $b_1(M_2) \geq 2$.*

Let $S^1 \rightarrow M_1$ be an essential circle in M_1 , and $S^1 \rightarrow M_2$ be a circle in M_2 with nonzero image in $H_1(M_2; \mathbb{Q})$. Then the $\mathbb{Z}$ -amalgamation

$$\pi_1(M_1 \cup_{S^1} M_2)$$

is not LERF.

Proof. At first, $b_1(M_2) \geq 2$ implies that, for any fiber bundle over circle structure $M_2 = N \times I / \phi$, $b_1(N) \geq 1$ holds. So any such N satisfies that $\pi_1(N)$ has a nontrivial finite quotient.

Take any fibered structure of M_2 , which gives $M_2 = N \times I / \phi$. By perturbing the fibered structure on M_2 , we can assume that $[N]$ has non zero algebraic intersection number with $[S^1] \in H_1(M_2; \mathbb{Z})$. Since $b_1(N) \geq 1$, $\pi_1(N)$ admits a finite quotient. So we are in the situation of Theorem 4.12, and $\pi_1(M_1 \cup_{S^1} M_2)$ is not LERF. $\square$

5. NONLERFNESS OF ARITHMETIC HYPERBOLIC MANIFOLD GROUPS

In this section, we will give the proof of Theorem 1.1 and Theorem 1.2, and give some results on nonLERFness of high dimensional nonarithmetic hyperbolic manifold groups. These results imply that most of known examples of high dimensional hyperbolic manifolds have nonLERF fundamental groups.

For all the proofs in this section, to prove a group is not LERF, we just show that it contains a subgroup isomorphic to one of the nonLERF groups in Theorem 1.3 or Theorem 1.4.

We start with the proof of Theorem 1.2, which claims that all noncompact arithmetic hyperbolic manifolds with dimension ≥ 4 have nonLERF fundamental groups.

Proof. We first consider the case that M is a noncompact standard arithmetic hyperbolic manifold.

We claim that M contains a (immersed) noncompact totally geodesic 3-dimensional submanifold N . This statement is well-known for experts, but the author did not find a reference on it. So we give a short proof here, which just uses linear algebra.

Since M is noncompact, it is defined by $\mathbb{Q}$ and a nondegenerate quadratic form $f : \mathbb{Q}^{m+1} \rightarrow \mathbb{Q}$ with negative inertia index 1. Let the symmetric bilinear form defining f be denoted by $B(\cdot, \cdot)$.

Since M is not compact, f represents 0 nontrivially in $\mathbb{Q}^{m+1}$, thus there exists $\vec{w} \neq \vec{0} \in \mathbb{Q}^{m+1}$ such that $B(\vec{w}, \vec{w}) = f(\vec{w}) = 0$. Since f is nondegenerate, there exists $\vec{v} \neq 0 \in \mathbb{Q}^{m+1}$ such that $B(\vec{v}, \vec{w}) \neq 0$. Let $V = \text{span}_{\mathbb{Q}}(\vec{v}, \vec{w})$. Then it is easy to see that $V^\perp \cap V = \{\vec{0}\}$ and the restriction of $B(\cdot, \cdot)$ on $V^\perp$ is positive definite.

Let $(\vec{v}_1, \dots, \vec{v}_{m-1})$ be a $\mathbb{Q}$ -basis of $V^\perp$ such that $B(\vec{v}_i, \vec{v}_j) = \delta_{ij}$ for $i, j \in \{1, \dots, m-1\}$. Then $W = \text{span}_{\mathbb{Q}}(\vec{v}_1, \vec{v}_2, \vec{v}, \vec{w})$ is a 4-dimensional subspace of $\mathbb{Q}^{m+1}$, such that the restriction of f on W has negative inertia index 1, and f represents 0 nontrivially on W .

So W and $f|_W$ define a (immersed) noncompact totally geodesic 3-dimensional suborbifold in $\mathbb{H}^m / SO_0(f, \mathbb{Z})$, which gives a (immersed) noncompact totally geodesic 3-dimensional submanifold N in M .

Now we are ready to prove the theorem. Here we consider M and N as compact manifolds, by adding Euclidean boundaries to the cusps.

Each boundary component of M has an Euclidean structure, so it is finitely covered by T^{m-1} . Each boundary component of N is homeomorphic to T^2 . We first take two copies of N . For each T^2 component of ∂N , take a long enough immersed $T^2 \times I$ in the corresponding boundary component of M , which is finitely covered by $T^{m-1} = (T^2 \times S^1) \times T^{m-4}$, such that the I factor wraps around the S^1 factor. This construction is same with the Freedman tubing construction in dimension 3. In [LR], it is shown that as long as the I factor wraps around the S^1 factor sufficiently many times, this immersed $N \cup (\partial N \times I) \cup N$ is π_1 -injective, so $\pi_1(N \cup (\partial N \times I) \cup N) < \pi_1(M)$.

Topologically, $N \cup (\partial N \times I) \cup N$ is just the double of N along ∂N . Since the double of N is a closed mixed 3-manifold with nontrivial geometric decomposition, Theorem 1.3 implies that $\pi_1(N \cup (\partial N \times I) \cup N)$ is not LERF. Then Lemma 2.3 implies $\pi_1(M)$ is not LERF.

Moreover, there exist both a free subgroup and a closed surface subgroup as a nonseparable subgroup in $\pi_1(M)$.

If M is a noncompact arithmetic hyperbolic manifold defined by quaternions, it also contains noncompact 3-dimensional totally geodesic submanifolds, by doing the same process for quadratic forms over quaternions. So the above proof also works in this case.

Since 7-dimensional arithmetic hyperbolic manifolds defined by octonions are all compact, the proof is done. $\square$

Then we give the proof of Theorem 1.1, which claims that all arithmetic hyperbolic manifolds with dimension ≥ 5 which are not those sporadic examples in dimension 7 have nonLERF fundamental groups. In this proof, we use two totally geodesic 3-dimensional submanifolds, instead of just using one such submanifold in the proof of Theorem 1.2.

Proof. We first suppose that M^m is a standard arithmetic hyperbolic manifold, with $m \geq 5$.

By the definition of standard arithmetic hyperbolic manifolds, there exists a totally real number field K , and a nondegenerate quadratic form $f : K^{m+1} \rightarrow K$ defined over K , such that the negative inertial index of f is 1 and f^σ is positive definite for all non-identity embeddings $\sigma : K \rightarrow \mathbb{R}$. Moreover, $\pi_1(M)$ is commensurable with $SO_0(f; \mathcal{O}_K)$. So to prove $\pi_1(M)$ is not LERF, we need only to show $SO_0(f; \mathcal{O}_K)$ is not LERF.

We can diagonalize the quadratic form f such that the symmetric matrix defining f is $A = \text{diag}(k_1, \dots, k_m, k_{m+1})$ with $k_1, \dots, k_m > 0$ and $k_{m+1} < 0$.

First suppose that there exists $i \in \{1, \dots, m\}$ such that $-\frac{k_i}{k_{m+1}}$ is not a square in K , and we can assume $i = 1$. Then f has two quadratic subforms defined by $\text{diag}(k_1, k_2, k_3, k_{m+1})$ and $\text{diag}(k_1, k_4, k_5, k_{m+1})$ respectively. These two subforms satisfy the conditions for defining arithmetic groups in $\text{Isom}_+(\mathbb{H}^3)$, and we denote these two subforms by f_1 and f_2 .

Then $SO_0(f_1; \mathcal{O}_K)$ and $SO_0(f_2; \mathcal{O}_K)$ are both subgroups of $SO_0(f; \mathcal{O}_K)$. Each of them fix a 3-dimensional totally geodesic plane in $\mathbb{H}^m$, and these two planes perpendicularly intersect with each other along a 1-dimensional biinfinite geodesic. (Here we do use that $m \geq 5$.) We denote these two 3-dimensional planes by P_1 and P_2 with $P_1 \cap P_2 = L$. Then $M_i = P_i / SO_0(f_i; \mathcal{O}_K)$ is a hyperbolic 3-orbifold for $i = 1, 2$. Moreover, it is easy to see that $SO_0(f_1; \mathcal{O}_K) \cap SO_0(f_2; \mathcal{O}_K) = SO_0(f_3; \mathcal{O}_K)$, with f_3 defined by $\text{diag}(k_1, k_{m+1})$ and $SO_0(f_3; \mathcal{O}_K)$ fixes L . The condition that $-\frac{k_1}{k_{m+1}}$ is not a square in K implies that f_3 does not represents 0 nontrivially in K^2 , so $SO_0(f_3; \mathcal{O}_K) \cong \mathbb{Z}$.

By a routine argument in hyperbolic geometry and using LERFness of hyperbolic 3-manifold groups (e.g. see Lemma 7.1 of [BHW]), there exist torsion free finite index subgroups $\Lambda_i < SO_0(f_i; \mathcal{O}_K)$ with $SO_0(f_3; \mathcal{O}_K) < \Lambda_i$ for $i = 1, 2$, and the subgroup of $SO_0(f; \mathcal{O}_K)$ generated by Λ_1 and Λ_2 is isomorphic to $\Lambda_1 *_\mathbb{Z} \Lambda_2$.

So $SO_0(f; \mathcal{O}_K)$ contains a subgroup $\Lambda_1 *_\mathbb{Z} \Lambda_2$, which is the fundamental group of $M_1 \cup_\gamma M_2$ for two hyperbolic 3-manifolds. By Theorem 1.4, $SO_0(f; \mathcal{O}_K)$ is not LERF, so $\pi_1(M)$ is also not LERF. Actually, we can also have that $\Lambda_1 *_\mathbb{Z} \Lambda_2 < \pi_1(M)$.

The nonseparable subgroup in $\pi_1(M)$ can be chosen to be a free product of surface groups and free groups. Moreover, if M is closed, then both M_1 and M_2 are closed, and the nonseparable subgroup can be chosen to be a free product of closed surface groups and free groups.

If $-\frac{k_i}{k_{m+1}}$ is a square in K for all $i \in \{1, \dots, m\}$, then the quadratic form f is equivalent to the diagonal form $\text{diag}(\underbrace{1, \dots, 1}_m, -1)$. It is easy to see that f is

also equivalent to the diagonal form $\text{diag}(2, 2, \underbrace{1, \dots, 1}_{m-2}, -1)$. So we reduce to the previous case.

If M^m is an arithmetic hyperbolic manifold defined by a quadratic form over quaternions, we can also find two totally geodesic 3-dimensional submanifolds intersecting along one circle. This can be done by diagonalizing the (skew-Hermitian) matrix with quaternion entries, then take two 2×2 submatrices with one common entry which contributes to the negative inertia index. Then the same proof as above also works here. $\square$

In [GPS], [Ag1] and [BT], the authors do cut-and-past surgery on standard arithmetic hyperbolic manifolds along codimension-1 totally geodesic arithmetic submanifolds, and constructed many nonarithmetic hyperbolic manifolds.

In [GPS], the authors took two non-commensurable standard arithmetic hyperbolic m -manifolds, cut them along isometric codimension-1 totally geodesic submanifolds, then glue them together by another way. This process is called "interbreeding". In [Ag1] and [BT], the authors cut one standard arithmetic hyperbolic m -manifold along two isometric codimension-1 totally geodesic submanifolds (the existence requires some knowledge in geometric group theory), then glue it back in a different way. This process is called "inbreeding", which is first carried out in [Ag1] for 4-dimensional case, and then generalized to higher dimensions in [BT].

Since all the resulted manifolds in the construction of [GPS], [Ag1] and [BT] contain codimension-1 totally geodesic arithmetic submanifolds, we have the following direct corollary of Theorem 1.1.

Theorem 5.1. *If M^m is a nonarithmetic hyperbolic m -manifold given by the constructions in [GPS] or [BT], with $m \geq 6$, then $\pi_1(M)$ is not LERF.*

Moreover, if M is closed, there exists a free product of closed surface groups and free groups as a nonseparable subgroup. If M is not closed, there exists either a free subgroup or a free product of closed surface groups and free groups as a nonseparable subgroup.

Another geometric and intuitive way of constructing hyperbolic m -manifolds is the reflection group method. Suppose P is a finite volume polyhedron in $\mathbb{H}^m$ such that any two codimension-1 faces which intersect with each other has dihedral angle $\frac{\pi}{n}$ with integer $n \geq 2$. Then the group generated by the reflections along codimension-1 faces of P is a discrete subgroup of $\text{Isom}(\mathbb{H}^m)$ with finite covolume.

For any torsion-free finite index subgroup of a reflection group consisting of orientation preserving isometries, then the corresponding quotient of $\mathbb{H}^m$ is a finite volume hyperbolic m -manifold M . M is a closed manifold if and only if P is compact in $\mathbb{H}^m$. The hyperbolic manifolds constructed by this method are not necessarily arithmetic, and it is known that there exist closed nonarithmetic reflection hyperbolic manifolds with dimension ≤ 5 , and noncompact nonarithmetic reflection hyperbolic manifolds with dimension ≤ 10 ([VS] Chapter 6.3.2).

In the case that $m \geq 5$, it is easy to see that M still contains two totally geodesic 3-dimensional submanifolds intersecting along a closed geodesic. To get such a picture, we take two totally geodesic 3-dimensional planes in $\mathbb{H}^m$ which contain two 3-dimensional faces of P and intersect with each other along one edge of P , then take their images in M . Similarly, for any $m \geq 4$, noncompact reflection hyperbolic m -manifolds also have noncompact totally geodesic 3-dimensional submanifolds, by taking the image of a 3-dimensional plane in $\mathbb{H}^m$ containing a noncompact 3-dimensional face of P .

So we get the following theorem for finite volume hyperbolic manifolds arising from reflection groups. The proof is exactly the same as the proof of Theorem 1.1 and Theorem 1.2.

Theorem 5.2. *Let M be a closed hyperbolic m -manifold such that $m \geq 5$, or a noncompact finite volume hyperbolic m -manifold with $m \geq 4$. If $\pi_1(M)$ is commensurable with the reflection group of some finite volume polyhedron in $\mathbb{H}^m$, then $\pi_1(M)$ is not LERF.*

Moreover, If M is closed, there exists a free product of closed surface groups and free groups as a nonseparable subgroup. If M is noncompact, there exists both a free subgroup and a closed surface subgroup as nonseparable subgroups.

By the dimension reason, there are no π_1 -injective $M_1 \cup_\gamma M_2$ submanifold in a 4-dimensional (arithmetic) hyperbolic manifold, so Theorem 1.4 does not give us any nonLERF fundamental group in dimension 4.

6. FURTHER QUESTIONS

In this section, we raise a few questions related to the results in this paper.

1. In Remark 3.9, we get that, for any mixed 3-manifold M , there exist a finite cover M' of M and a π_1 -injective properly immersed subsurface $\Sigma \looparrowright M'$, such that $\pi_1(\Sigma)$ is not contained in any finite index subgroup of $\pi_1(M')$. We may ask whether taking this finite cover is necessary.

Question 6.1. For any mixed 3-manifold M , whether there exists a π_1 -injective (properly) immersed subsurface $\Sigma \looparrowright M$, such that $\pi_1(\Sigma)$ is not contained in any finite index subgroup of $\pi_1(M)$?

2. None of the results in this paper cover the case of compact (arithmetic) hyperbolic 4-manifolds, since they neither contain $M_1 \cup_\gamma M_2$ as a singular submanifold, nor contain a $\mathbb{Z}^2$ subgroup (and a mixed 3-manifold group as a subgroup).

One possible approach for compact (arithmetic) hyperbolic 4-manifolds is to study the group of $M_1 \cup_S M_2$ with M_1 and M_2 being compact arithmetic hyperbolic 3-manifolds, with S being a hyperbolic surface embedded in both M_1 and M_2 . In this case, the edge group is a closed surface group, which is much more complicated than $\mathbb{Z}$ or $\mathbb{Z}^2$. The method in this paper seems do not work directly in this case. Even if it works (under some clever modification), the nonseparable (finitely generated) subgroup constructed by this method would be infinitely presented.

3. Given the nonLERFness results of high dimensional (arithmetic) hyperbolic manifolds in this paper, maybe it is not too ambitious to ask the following question about general high dimensional hyperbolic manifolds.

Question 6.2. Whether all finite volume hyperbolic manifolds with dimension at least 4 have nonLERF fundamental groups?

As mentioned previously, the 4-dimensional compact case is tricky. Another problem is that we do not have many examples of finite volume high dimensional hyperbolic manifolds. To the best of the authors knowledge, the only methods for constructing high dimensional hyperbolic manifolds (dimension ≥ 4) are: the arithmetic method, the cut-and-paste method and the reflection group method. In this paper, it is shown in Theorem 1.1 and Theorem 1.2, Theorem 5.1, and Theorem 5.2 that these three constructions give nonLERF fundamental groups in dimension ≥ 5 (not covering 7-dimensional sporadic examples), ≥ 6 and ≥ 5 respectively.

However, it is difficult to understand a general high dimensional hyperbolic manifold, if we do not assume it lies in one of the above families. The author does not know whether a general high dimensional hyperbolic manifold group contains 3-manifold subgroups. Maybe a generalization of [KM] (which shows that each closed hyperbolic 3-manifold admits a π_1 -injective immersed almost totally geodesic closed subsurface) can do this job, but it seems very difficult for the author.

4. The author expects the method in this paper can be used to prove more groups are not LERF. However, since the author does not have very broad knowledge in group theory, we only consider the groups of finite volume hyperbolic manifolds in this paper, which is one of the author's favorite family of groups.

The author also expects that the method in this paper can be translated to a purely algebraic proof, instead of a geometric one. Actually, most part of the proof are essentially algebraic, except for one point. In Proposition 3.6 and Proposition 3.7 (also Proposition 4.8 and Proposition 4.10), although the essential part which gives the nonseparability is $\Sigma_{1,1} \cup \Sigma_{2,1}$, we still need to take a bigger (singular) surface so that it defines a nontrivial 1-dimensional cohomology class in some finite cover. Then we take a proper cyclic finite cover dual to this cohomology class and get a contradiction. Although this process seems can be done algebraically, the author does not know how to work it out.

REFERENCES

- [Ag1] I. Agol, *Systoles of hyperbolic 4-manifolds*, preprint 2006, <http://arxiv.org/abs/math/0612290>.
- [Ag2] I. Agol, *The virtual Haken conjecture*, with an appendix by I. Agol, D. Groves, J. Manning, Documenta Math. 18 (2013), 1045 - 1087.
- [BT] M. Belolipetsky, S. Thomson, *Systoles of hyperbolic manifolds*, Algebr. Geom. Topol. 11 (2011), no. 3, 1455 - 1469.
- [BHW] N. Bergeron, F. Haglund, D. Wise, *Hyperplane sections in arithmetic hyperbolic manifolds*, J. Lond. Math. Soc. (2) 83 (2011), no. 2, 431 - 448.
- [BBS] A. Brunner, R. Burns, D. Solitar, *The subgroup separability of free products of two free groups with cyclic amalgamation*, Contributions to group theory, 90 - 115, Contemp. Math., 33, Amer. Math. Soc., Providence, RI, 1984.
- [CDW] E. Chesebro, J. DeBlois, H. Wilton, *Some virtually special hyperbolic 3-manifold groups*, Comment. Math. Helv. 87 (2012), no. 3, 727 - 787.
- [DLW] P. Derbez, Y. Liu, S. Wang, *Chern-Simons theory, surface separability, and volumes of 3-manifolds*, J. Topol. 8 (2015), no. 4, 933 - 974.
- [Gi] R. Gitik, *Graphs and separability properties of groups*, J. Algebra 188 (1997), no. 1, 125 - 143.
- [GPS] M. Gromov, I. Piatetski-Shapiro, *Nonarithmetic groups in Lobachevsky spaces*, Inst. Hautes Études Sci. Publ. Math. No. 66 (1988), 93 - 103.
- [Ha] M. Hall, *Coset representations in free groups*, Trans. Amer. Math. Soc. 67 (1949), 421 - 432.
- [KM] J. Kahn, V. Markovic, *Immersing almost geodesic surfaces in a closed hyperbolic three manifold*, Ann. of Math. (2) 175 (2012), no. 3, 1127 - 1190.
- [Ka] M. Kapovich, *Noncoherence of arithmetic hyperbolic lattices*, Geom. Topol. 17 (2013), no. 1, 39 - 71.
- [LM] J. Li, J. Millson, *On the first Betti number of a hyperbolic manifold with an arithmetic fundamental group*, Duke Math. J. 71 (1993), no. 2, 365 - 401.
- [Li1] Y. Liu, *A characterization of virtually embedded subsurfaces in 3-manifolds*, Trans. Amer. Math. Soc., to appear, preprint 2014, <http://arxiv.org/abs/1406.4674>.
- [Li2] Y. Liu, *Erratum to "A characterization of virtually embedded subsurfaces in 3-manifolds"*, preprint 2016.
- [LR] D. Long, A. Reid, *The fundamental group of the double of the figure-eight knot exterior is GFERF*, Bull. London Math. Soc. 33 (2001), no. 4, 391 - 396.
- [Ni1] G. Niblo, *The subgroup separability of amalgamated free products*, Ph.D. Thesis, University of Liverpool, 1988.
- [Ni2] G. Niblo, *H.N.N. extensions of a free group by $\mathbb{Z}$ which are subgroup separable*, Proc. London Math. Soc. (3) 61 (1990), no. 1, 18 - 32.
- [NW1] G. Niblo, D. Wise, *The engulfing property for 3-manifolds*, The Epstein birthday schrift, 413 - 418 (electronic), Geom. Topol. Monogr., 1, Geom. Topol. Publ., Coventry, 1998.

- [NW2] G. Niblo, D. Wise, *Subgroup separability, knot groups and graph manifolds*, Proc. Amer. Math. Soc. 129 (2001), no. 3, 685 - 693.
- [PW1] P. Przytycki, D. Wise, *Mixed 3-manifolds are virtually special*, preprint 2012, <http://arxiv.org/abs/1205.6742>.
- [PW2] P. Przytycki, D. Wise, *Separability of embedded surfaces in 3-manifolds*, Compos. Math. 150 (2014), no. 9, 1623 - 1630.
- [Ri] E. Rips, *An example of a non-LERF group which is a free product of LERF groups with an amalgamated cyclic subgroup*, Israel J. Math. 70 (1990), no. 1, 104 - 110.
- [RW] H. Rubinstein, S. Wang, *π_1 -injective surfaces in graph manifolds*, Comment. Math. Helv. 73 (1998), no. 4, 499 - 515.
- [Sc] P. Scott, *Subgroups of surface groups are almost geometric*, J. London Math. Soc. (2) 17 (1978), no. 3, 555 - 565.
- [Th] W. Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. 59 (1986), no. 339, i - vi and 99 - 130.
- [VS] E. Vinberg, O. Shvartsman, *Discrete groups of motions of spaces of constant curvature*, Geometry, II, 139 - 248, Encyclopaedia Math. Sci., 29, Springer, Berlin, 1993.
- [Wi] D. Wise, *The structure of groups with a quasiconvex hierarchy*, preprint, <http://www.math.mcgill.ca/wise/papers.html>.

DEPARTMENT OF MATHEMATICS, UC BERKELEY, CA 94720, USA
E-mail address: hongbins@math.berkeley.edu